arXiv:1106.2341v2 [math.GT] 21 Aug 2011

Linearly Bounded Conjugator Property for Mapping Class Groups

Jing Tao

Abstract

Given two conjugate mapping classes f and g, we produce a conjugating element ω such that $|\omega| \leq K\big(|f|+|g|\big)$, where $|\cdot|$ denotes the word metric with respect to a fixed generating set, and K is a constant depending only on the generating set. As a consequence, the conjugacy problem for mapping class groups is exponentially bounded.

1 Introduction

Two fundamental problems in group theory posed by Dehn are the word problem and the conjugacy problem [Deh11]. Given a group with a fixed presentation, the *word problem* asks if there is an algorithm that can decide in finite time if a given word is the identity. The *conjugacy problem* seeks an algorithm to decide if two words represent the same conjugacy class. Since the conjugacy class of the identity element is itself, the word problem can be seen as a special case of the conjugacy problem. Not all groups have solvable word problem [Boo59], hence the same is true for the conjugacy problem.

In this paper, we are interested in these problems for mapping class groups $\mathcal{MCG}(\mathcal{S})$ of surfaces $\mathcal{S}$ of finite type. We establish the following:

***Theorem* A.** *There is an exponential-time algorithm to solve the conjugacy problem for* $\mathcal{MCG}(\mathcal{S})$.

There is some history to the word and conjugacy problems for $\mathcal{MCG}(\mathcal{S})$. The first solution to the word problem can be attributed to Grossman, whose actual contribution is proving residual finiteness for $\mathcal{MCG}(\mathcal{S})$ [Gro75]. In [Mos95], Mosher showed $\mathcal{MCG}(\mathcal{S})$ admits an automatic structure, from which a quadratic-time solution to the word problem is obtained. (See [ECH$^+$92] for a background on automatic groups). It is not yet known if a sub-quadratic solution is possible. In [Hem79], Hemion solved the conjugacy problem for $\mathcal{MCG}(\mathcal{S})$, but his algorithm is not exponentially bounded. In [Mos86], Mosher gave a faster algorithm for deciding conjugacy among pseudo-Anosov mapping classes. (A similar result was recently obtained by Agol in [Ago10].) Using the work of Bestvina-Handel [BH95], which gives an algorithm for detecting pseudo-Anosovs, Mosher extended his result to compute complete conjugacy invariants for all mapping classes [Mos03].

Our strategy to Theorem A is to apply Mosher's automaticity result. In general, a solution to the word problem does not necessarily yield a solution to the conjugacy problem; it is an open question whether all automatic groups have solvable conjugacy problem [ECH$^+$92]. A sufficient condition is if the group has *linearly bounded conjugator (L.B.C.) property* (see theorem below or Definition 2.2.1). The main theorem of our paper is that L.B.C. property is satisfied by $\mathcal{MCG}(\mathcal{S})$.

***Theorem* B** (L.B.C. property for $\mathcal{MCG}(\mathcal{S})$)**.** *Let Λ be a finite generating set for $\mathcal{MCG}(\mathcal{S})$. There exists a constant K, depending only on Λ, such that if $f, g \in \mathcal{MCG}(\mathcal{S})$ are conjugate, then there is a conjugating element ω with*

$$|\omega| \leq K\big(|f| + |g|\big).$$

To see how Theorem A follows from Theorem B, we give an algorithm to the conjugacy problem. Given two arbitrary elements $f, g \in \mathcal{MCG}(\mathcal{S})$, let B be the ball of radius $K\big(|f|+|g|\big)$ in $\mathcal{MCG}(\mathcal{S})$. To decide if f and g are conjugate it suffices to check if $\omega \in B$ satisfies $\omega f \omega^{-1} g^{-1} = 1$. We run Mosher's quadratic algorithm to the word problem to all words of the form $\omega f \omega^{-1} g^{-1}$ with $\omega \in B$. The number of elements in B is an exponential function of the radius, therefore the complexity of this solution is an exponential function of the word lengths of f and g.

Linearly-bounded conjugator property is satisfied by hyperbolic groups [Lys89, Lemma 10], as well as by torsion elements in groups acting on CAT(0) spaces [BH99, III.1.13]. These are important classes of groups which have solvable word and conjugacy problems [Gro87, BH99]. Hyperbolic groups in fact have efficient algorithms: the word problem is solvable in linear time, and the conjugacy problem in quadratic time [BH99]. As long as the surface $\mathcal{S}$ has disjoint isotopy classes of curves, $\mathcal{MCG}(\mathcal{S})$ is not hyperbolic, as Dehn twists about disjoint curves give rise to higher rank free abelian subgroups. It is also known that $\mathcal{MCG}(\mathcal{S})$ does not act on any complete CAT(0) space [BH99, II.7.26]. Nevertheless, $\mathcal{MCG}(\mathcal{S})$ shares many properties with hyperbolic groups, and much of the pursuit in its study has been to understand to what extent it resembles and differs from hyperbolic groups. Establishing L.B.C. property for $\mathcal{MCG}(\mathcal{S})$ thus provides another positive analogy between $\mathcal{MCG}(\mathcal{S})$ and hyperbolic groups.

After we announced our result, Hamenstädt in [Ham09] announced biautomaticity for $\mathcal{MCG}(\mathcal{S})$, which generalizes Mosher's automaticity result as well as obtains Theorem A. Another consequence of her work is the exponentially-bounded conjugator property for $\mathcal{MCG}(\mathcal{S})$. Notice, however, that this bound only gives a doubly-exponential solution to the conjugacy problem if we use the same algorithm as described below Theorem B, since the search space for the conjugator would grow doubly-exponential in terms of the word lengths of the elements.

1.1 Idea of the proof of Theorem B

The proof of Theorem B is broken up into three arguments, following the classification of mapping classes into pseudo-Anosov, reducible, and finite order. The case of pseudo-Anosovs was settled by Masur-Minsky in [MM00], using the machinery of hierarchies developed in the same paper. This paper resolves the other two cases. Surprisingly, it turns out that the most delicate case involves the finite order elements of $\mathcal{MCG}(\mathcal{S})$. In many ways, pseudo-Anosov mapping classes can be viewed as the "hyperbolic" elements of $\mathcal{MCG}(\mathcal{S})$, whereas the finite order mapping classes are the "elliptic" ones. The methods that Masur and Minsky developed are suited for elements that behave more hyperbolically, and thus are not effective for finite order mapping classes. Our main contribution is the development of new tools for the study of finite order mapping classes. Just as in the case of pseudo-Anosov mapping classes, we rely heavily on the machinery of hierarchies, which we need to extend so it is more suited to deal with the elliptic geometry.

We briefly explain how hierarchies are related to words in $\mathcal{MCG}(\mathcal{S})$. A natural model space for $\mathcal{MCG}(\mathcal{S})$ is the marking graph $\mathrm{Mark}(\mathcal{S})$ of $\mathcal{S}$. A marking $\mu_0 \in \mathrm{Mark}(\mathcal{S})$ is a collection of curves on $\mathcal{S}$ satisfying certain technical conditions. Given an element $f \in \mathcal{MCG}(\mathcal{S})$, the image of μ_0 under f determines f up to finitely many choices. Paths from μ_0 to $f\mu_0$ are naturally associated to words representing f. Even though $\mathrm{Mark}(\mathcal{S})$ (or any other model space) is not hyperbolic or CAT(0), there is a coarsely well-defined projection map $\mathrm{Mark}(\mathcal{S})$ to an infinite product of hyperbolic spaces. These hyperbolic spaces are the curve complexes of essential subsurfaces (including annuli) of $\mathcal{S}$ [MM99]. The projection of μ_0 and $f\mu_0$ to each curve complex gives rise a family of paths in curve complexes. Hierarchies organize these paths to become preferred quasi-geodesics in $\mathrm{Mark}(\mathcal{S})$ [MM00].

The hyperbolic geometry of pseudo-Anosovs are exhibited in the fact that they act hyperbolically (with north-south dynamics) on curve complexes [MM99, MM00]. This is analogous to the way how the infinite order elements of a hyperbolic group act on the Cayley graph of the group.

Hierarchies are used to build quasi-axes in curve complexes for the action of pseudo-Anosovs. Then, Masur and Minsky proved a fellow-traveling type property for hierarchies which can be applied to fellow-traveling quasi-axes. This allowed them to extend the proof of L.B.C. property from infinite-order elements of hyperbolic groups to pseudo-Anosovs [MM00, Theorem 7.2].

The appropriate analogy for finite order mapping classes are the torsion elements of a group G that acts on a CAT(0) (or hyperbolic) space X. We are inspired by the argument contained in [BH99] on L.B.C. property for torsion elements of G which we will briefly sketch. In our definition, an element $g \in G$ acts *elliptically* on X if it satisfies two conditions. First, g acts on X with (coarse) fixed points. Second, every orbit of g has a center of mass whose distance to the orbit is comparable to the word length of g. Since X is CAT(0) (or hyperbolic), g is torsion implies g is elliptic. After conjugating by an appropriate element, the center of mass can be moved into a fixed ball containing the fundamental domain for the action of G on X. The set of torsion elements of G having fixed points inside the ball is finite and contains a representative for each conjugacy class. From here, one can reduce L.B.C. property in the elliptic case to a finite set.

We want to show finite order mapping classes act elliptically on $\mathrm{Mark}(\mathcal{S})$. Since curve complexes are hyperbolic, the action of a finite order element $f \in \mathcal{MCG}(\mathcal{S})$ is elliptic on the curve complex $\mathcal{C}(Z)$ of Z, for any essential subsurface $Z \subseteq \mathcal{S}$. Let $\pi_Z : \mathrm{Mark}(\mathcal{S}) \to \mathcal{C}(Z)$ be the projection map. The orbit of $\mu_0 \in \mathrm{Mark}(\mathcal{S})$ under f maps uniformly close to the orbit of $\pi_Z(\mu_0)$ under f in $\mathcal{C}(Z)$. Philosophically, what we want is a marking $\mu \in \mathrm{Mark}(\mathcal{S})$ such that, for all Z, $\pi_Z(\mu)$ lies uniformly close to the center of mass of the orbit of $\mu_Z(\mu_0)$. If such a μ exists, then by the distance formula (Theorem 2.6.5), μ is a coarse fixed point of f in $\mathrm{Mark}(\mathcal{S})$ with uniform coarse constant. However, knowing that $\pi_Z(\mu)$ is close to the center of mass in each $\mathcal{C}(Z)$ is not enough to guarantee the second condition of elliptic action for f on $\mathrm{Mark}(\mathcal{S})$. We have to know the exact projection of μ to $\mathcal{S}$ and to families of subsurfaces of $\mathcal{S}$ which are pairwise disjoint. (This is the reason we do not approach this problem using the Consistency theorem of [BKMM06].) Our approach is a balance act between finding a marking which is close enough to the orbit to satisfy the second condition while controlling the coarse constant so that the marking satisfies the first condition for some constant. Our method is constructive and utilizes the symmetry of the action of f on $\mathcal{S}$. We prove

***Theorem* C.** *Finite order mapping classes elliptically act on* $\mathrm{Mark}(\mathcal{S})$. *More precisely, given* $\mu_0 \in \mathcal{MCG}(\mathcal{S})$. *There exists a constant* k *depending only on* μ_0 *such that, for any finite order element* $f \in \mathcal{MCG}(\mathcal{S})$, *there exists a marking* $\mu \in \mathrm{Mark}(\mathcal{S})$ *coarsely fixed by* f *by a uniform constant satisfying*

$$d_{\mathrm{Mark}(\mathcal{S})}(\mu_0, \mu) \leq k|f|.$$

***Corollary* D.** *There exist a constant* K, *depending only on* $\mathcal{S}$, *and a finite set of elements* $\Sigma \subset \mathcal{MCG}(\mathcal{S})$ *such that if* $f \in \mathcal{MCG}(\mathcal{S})$ *has finite order, then there exists* $\omega \in \mathcal{MCG}(\mathcal{S})$ *such that* $\omega f \omega^{-1} \in \Sigma$ *and* $|\omega| \leq K|f|$.

For reducible mapping classes, we combine the two arguments above. If f is a reducible mapping class of infinite order, then up to taking powers the surface $\mathcal{S}$ can be decomposed into a collection of subsurfaces on which f is either pseudo-Anosov or has finite order. In order to apply induction to subsurfaces, we need to built paths from μ_0 to $f\mu_0$ in $\mathrm{Mark}(\mathcal{S})$ that move only in the complementary subsurfaces of the reducing system of f. This is possible if the initial marking μ_0 contains the reducing system of f. However, one marking cannot contain all possible reducing systems, even up to conjugation. But it suffices to reduce to a finite problem. We prove

***Theorem* E.** *There exist a constant* K *and a finite set of markings* $\mathcal{M}$ *so that if* $f \in \mathcal{MCG}(\mathcal{S})$ *is reducible, then there exists* $\omega \in \mathcal{MCG}(\mathcal{S})$ *such that the reducing system of* $\omega f \omega^{-1}$ *is contained in some* $\mu \in \mathcal{M}$ *and* $|\omega| \leq K|f|$.

Finally, each case in the classification will produce a different constant. The proof of L.B.C. property for $\mathcal{MCG}(\mathcal{S})$ will be complete by taking a maximum over the three constants.

The organization of the paper is as follows.

- In §2, we review basic definitions and the theory of hierarchies. A key notion that we will introduce is the notion of *separating markings* in §2.9.
- In §3, we give a couple of definitions and prove two technical lemmas about finite order mapping classes which will be useful for the next section. We also construct an example which motivates this section and the next section.
- In §4, we prove Theorem C and derive L.B.C. property for finite order mapping classes.
- In §5, we prove Theorem E and use the known results for pseudo-Anosov and finite order elements to derive L.B.C. property for infinite order reducible mapping classes.

1.2 Acknowledgments

The author would like to thank her thesis advisor, Howard Masur, for his excellent guidance and for suggesting the problem. She would also like to thank Kasra Rafi for helpful conversations throughout this project, and also Benson Farb, Daniel Groves, and Chris Leininger for helpful suggestions and their interest in this work.

2 Preliminaries

In this section, we develop the background material for the paper. Our main tool will be Masur and Minsky's theory of hierarchies. From §2.6 to §2.8, we will summarize the properties of hierarchies that will be needed for this paper. Some of the definitions will be merely sketched and most of the proofs will be omitted. We refer the reader to Masur-Minsky's paper [MM00] for more details. We also refer to [FLP79] and [FM08] for general references on mapping class groups and the topology of surfaces, and to [Gro87] and [BH99] for references on δ-hyperbolic spaces.

2.1 Arcs, curves and surfaces

Let $\mathcal{S} = \mathcal{S}_{g,p}$ be a connected, oriented surface of genus g with p punctures. We call $\xi(\mathcal{S}) = 3g-3+p$ the *complexity* of $\mathcal{S}$. Surfaces of complexity strictly greater than 1 are called *generic* surfaces. Surfaces of complexity 1 are called *sporadic* and they are topologically either the four-holed sphere or one-holed torus. Two remaining low-complexity cases are *exceptional surfaces.* Complexity 0 is the three-holed sphere or a *pair of pants*, and complexity -1 is topologically an annulus. Throughout this paper we will be working with a generic surface $\mathcal{S}$ without boundary. But sporadic and exceptional surfaces and surfaces with boundary naturally arise as subsurfaces of $\mathcal{S}$, and thus are important for induction arguments.

An *essential curve* or just a *curve* on $\mathcal{S}$ will always mean the free isotopy class of a simple closed curve, which is not null-homotopic or homotopic to a puncture or a boundary component. The *geometric intersection number* $i(\alpha, \beta)$ of a pair of curves α and β is the minimal number of intersections among representatives of α and β. A *multicurve* or a *curve system* will mean a finite collection of distinct curves which can be realized disjointly. A *pants decomposition* of $\mathcal{S}$ is a maximal curve system c on $\mathcal{S}$. In particular, each component of $\mathcal{S} \setminus c$ is topologically a pair of pants. Note that a pants decomposition exists for $\mathcal{S}$ if only if $\xi(\mathcal{S}) \geq 1$, in which case the cardinality of c is equal to $\xi(\mathcal{S})$. To talk about arcs we need $\mathcal{S}$ to have boundary. An *arc* will be an isotopy class of a simple arc δ, with isotopies relative to the boundary, such that δ has both endpoints on $\partial\mathcal{S}$ and is not isotopic to a boundary component. The geometric intersection number between two arcs will be the number of intersections in the interior of $\mathcal{S}$ modulo isotopies relative to the boundary of $\mathcal{S}$.

A *subsurface* Y of $\mathcal{S}$ is an isotopy class of a connected and open subsurface $Y \subset \mathcal{S}$ such that $\partial\overline{Y}$ are essential curves. We include the possibility that $Y = \mathcal{S}$ unless we say a *proper* subsurface. An annular subsurface A of $\mathcal{S}$ is a regular neighborhood of a curve α with simple boundaries. We will often abuse terminology by confusing A with its *core curve* α, and refer to α as a subsurface of $\mathcal{S}$ as well. In this case, we will let $\partial A = \alpha$. For reasons we shall see, we will distinguish subsurfaces which are not pants, called *essential subsurfaces* or *domains*.

To do away with isotopy classes, we endow $\mathcal{S}$ with a hyperbolic metric, so any curve α has a unique geodesic representative α^*, and $i(\alpha, \beta) = i(\alpha^*, \beta^*)$. Each curve system can be assumed to consist of geodesic representatives. Similarly, each non-annular subsurface of $\mathcal{S}$ also has a unique representative.

2.2 Mapping class groups

Let $\mathrm{Homeo}^+(\mathcal{S})$ be the group of orientation-preserving self-homeomorphisms of $\mathcal{S}$. The *mapping class group* of $\mathcal{S}$ is

$$\mathcal{MCG}(\mathcal{S}) = \mathrm{Homeo}^+(\mathcal{S})/\sim$$

where $f \sim g$ if and only if $g^{-1} \circ f$ is isotopic to the identity on $\mathcal{S}$. Elements of $\mathcal{MCG}(\mathcal{S})$ are called mapping classes. It is well-known that $\mathcal{MCG}(\mathcal{S})$ is finitely generated (and finitely presented) [Lic64]. For this paper, we will fix a finite generating set Λ of $\mathcal{MCG}(\mathcal{S})$. We will often regard $\mathcal{MCG}(\mathcal{S})$ as a metric space by considering the word metric $|\cdot| = |\cdot|_\lambda$ induced by Λ.

If $\mathcal{S}$ is a once-punctured torus or four-times punctured sphere, then $\mathcal{MCG}(\mathcal{S})$ is commensurable to $\mathrm{SL}(2, \mathbb{Z})$. The mapping class group of a thrice-punctured sphere is finite. For us, an annulus A will always appear as a regular neighborhood of a simple closed curve on an ambient surface, so A has two boundary components. Let $\mathcal{MCG}(A, \partial A)$ be the group of isotopy classes of homeomorphisms of A relative to ∂A. One checks that $\mathcal{MCG}(A, \partial A) \simeq \mathbb{Z}$.

Definition 2.2.1 (L.B.C. property)**.** Given a finitely generated group G equipped with a finite generating set Λ, we say a conjugacy class $\mathfrak{c}$ of G has *linearly bounded conjugators* if for any $f, g \in \mathfrak{c}$, there exists a conjugating element $\omega \in G$ such that

$$|\omega| \leq K_{\mathfrak{c}}\big(|f| + |g|\big),$$

where $|\cdot|$ represent the word length in Λ, and $K_{\mathfrak{c}}$ depends only on $\mathfrak{c}$ and Λ. If $K = K_{\mathfrak{c}}$ can be taken to be independent of the conjugacy class $\mathfrak{c}$, then we say G has *linearly bounded conjugator property* or *L.B.C. property.* If G has L.B.C. property for Λ, then changing Λ to any other finite generating set changes K by a bounded amount. Therefore, the definition is independent of the choice of the generating set, so Λ can always be taken to be a symmetric generating set.

Mapping class groups of non-generic surfaces satisfy L.B.C. property. We would like to show the same is true for mapping class groups of generic surfaces. The first observation is that the Nielsen-Thurston classification of mapping classes is a conjugacy invariant. This means that we can argue for L.B.C. property separately for each type. We refer to [Thu88] and [FM08, §13] for more details on the classification theorem. Recall that a mapping class f is called *irreducible* if f does not fix any multicurve (setwise); otherwise f is called *reducible.* The following statement applies to all surfaces $\mathcal{S}$.

Theorem 2.2.2 (Nielsen-Thurston classification for $\mathcal{MCG}(\mathcal{S})$)**.** *Every element $f \in \mathcal{MCG}(\mathcal{S})$ is either pseudo-Anosov, periodic (finite order), or reducible. Furthermore, for each $f \in \mathcal{MCG}(\mathcal{S})$, let σ be a (possibly empty) multicurve fixed by f. Let $Y_1, \ldots, Y_k$ be the connected components of $\mathcal{S} \setminus \sigma$, and for each i, choose the smallest $n_i \in \mathbb{N}$ so that $f^{n_i}(Y_i) = Y_i$. Then for any i, $f^{n_i}|_{Y_i}$ either has finite order or is pseudo-Anosov.*

The set of curves σ satisfying Theorem 2.2.2 f is called a *reducing system* for f. For each $Y_i \in \mathcal{S} \setminus \sigma$, the map f^{n_i} is called the *first return map of f to Y_i*. Note that the first return map of f to Y_i means exactly that $f^{n_i}|_{Y_i}$ can be viewed as an element of $\mathcal{MCG}(Y_i)$. The content of the classification can be rephrased to say f is pseudo-Anosov if and only if f is irreducible of infinite order.

Definition 2.2.3 (Canonical reducing systems)**.** By choosing σ to be a minimal collection of curves satisfying Theorem 2.2.2, then $\sigma = \sigma_f$ is unique up to isotopy and is called the *canonical reducing system* for f. If $f \in \mathcal{MCG}(\mathcal{S})$ is either pseudo-Anosov or finite order, then $\sigma_f = \emptyset$. (See [Mos07]).

In [MM00, §7], Masur-Minsky proved L.B.C. property for the pseudo-Anosov elements of $\mathcal{MCG}(\mathcal{S})$.

Theorem 2.2.4 (L.B.C. property for pseudo-Anosov mapping classes)**.** *There exists a constant K, depending only on $\mathcal{S}$, such that if $f, g \in \mathcal{MCG}(\mathcal{S})$ are conjugate pseudo-Anosov mapping classes, then there is a conjugating element $\omega \in \mathcal{MCG}(\mathcal{S})$ with*

$$|\omega| \leq K\big(|f| + |g|\big).$$

Our goal in this paper is to prove L.B.C. property for the finite order and reducible elements of $\mathcal{MCG}(\mathcal{S})$. The argument for finite order mapping classes is the hard part of this paper. The argument for reducible mapping classes is inductive and will make use of the canonical reducing system.

2.3 Complexes of curves

The *complex of curves* $\mathcal{C}(\mathcal{S})$ on a surface $\mathcal{S}$ is a locally-infinite, finite dimensional simplicial complex on which $\mathcal{MCG}(\mathcal{S})$ acts by automorphisms. The definition first appeared in [Har81]. We treat generic, sporadic, and exceptional surfaces separately.

- **Generic surfaces.** Suppose $\mathcal{S}$ has $\xi(\mathcal{S}) > 1$. The k–th skeleton $\mathcal{C}_k(\mathcal{S})$ consists of all curve systems on $\mathcal{S}$ of cardinality $k+1$. There is an obvious inclusion of $\mathcal{C}_{k-1}(\mathcal{S}) \hookrightarrow \mathcal{C}_k(s)$ by face relations. Top dimensional simplices of $\mathcal{C}(\mathcal{S})$ correspond to pants decompositions on $\mathcal{S}$, hence $\dim(\mathcal{C}(\mathcal{S})) = \xi(\mathcal{S}) - 1$.
- **Sporadic surfaces.** With the above definition, the curve complex of a sporadic surface $\mathcal{S}$ would be a disconnected set of points. To construct a more useful object, we modify the definition to allow two vertices in $\mathcal{C}(\mathcal{S})$ span an edge if they intersect minimally over $\mathcal{S}$ (once for one-holed torus and twice for four-holed sphere). It is a classical theorem that with this definition $\mathcal{C}(\mathcal{S})$ is isomorphic to the Farey graph [HT80, Min96].
- **Pants.** A pair of pants has no essential curves. Here we do not modify the definition and let the curve complex of pants be empty. This is the reason why we want to exclude pants from essential subsurfaces.
- **Annuli.** An arbitrary annulus has no essential curves. But for us, an annulus A will always appear as a regular neighborhood of a curve γ in a larger surface, and we would like $\mathcal{C}(A)$ (or $\mathcal{C}(\gamma)$) to record twist information about γ. Vertices of $\mathcal{C}(A)$ will be properly embedded arcs and two arcs are connected by an edge if they can be isotoped rel endpoints to have disjoint interiors.

By an element or subset of $\mathcal{C}(\mathcal{S})$ we will always mean an element or subset of $\mathcal{C}_0(\mathcal{S})$. We make $\mathcal{C}(\mathcal{S})$ into a complete geodesic metric space by endowing each simplex with an Euclidean structure with edge lengths 1. From the perspective of coarse geometry, we do not lose anything by identifying $\mathcal{C}(\mathcal{S})$ with its 1-skeleton. We denote by $d_{\mathcal{C}(\mathcal{S})}$, or more simply by $d_{\mathcal{S}}$, the shortest distance in $\mathcal{C}_1(\mathcal{S})$ between two vertices. If A is an annulus with a core curve γ, we will also use the notation d_γ or d_A to denote distances in $\mathcal{C}(A)$. Induction on intersection number can be used to show $\mathcal{C}(\mathcal{S})$ is connected, and $d_{\mathcal{S}}(\alpha,\beta) \leq i(\alpha,\beta)$. The simplicial action of $\mathcal{MCG}(\mathcal{S})$ on $\mathcal{C}(\mathcal{S})$ preserves this metric. The action is not proper. The quotient $\mathcal{C}(\mathcal{S})/\mathcal{MCG}(\mathcal{S})$ parametrizes curves on $\mathcal{S}$ up to homeomorphisms, hence it is finite.

For a generic surface $\mathcal{S}$, $d_{\mathcal{S}}$ coarsely measures the complexity between two curves in the following sense: $d_{\mathcal{S}}(\alpha,\beta) = 1$ if and only if α and β are disjoint; $d_{\mathcal{S}}(\alpha,\beta) = 2$ if and only if α and β cohabit a proper subsurface $Y \subset \mathcal{S}$; $d_{\mathcal{S}}(\alpha,\beta) \geq 3$ if and only α and β *fill* $\mathcal{S}$, or the complement of their union in $\mathcal{S}$ does not support any essential curve.

Paths in curve complexes are not unique. The following theorem in [MM99] gives us some geometric control over paths.

Theorem 2.3.1 ([MM99]). *For any surface $\mathcal{S}$ which is not a pair of pants, $\mathcal{C}(\mathcal{S})$ has infinite diameter and is δ–hyperbolic.*

For sporadic surfaces, Theorem 2.3.1 follows from a classical result that the Farey graph is quasi-isometric to an infinite-valence tree (see [Man05]). In the case of an annulus, Theorem 2.3.1 follows from the easy fact that $\mathcal{C}(A)$ is quasi-isometric to $\mathbb{Z}$.

For generic surfaces, there are several ways to see that $\mathcal{C}(\mathcal{S})$ has infinite diameter. Relevant to our paper is the following lemma.

Lemma 2.3.2 ([MM99, Proposition 4.6]). *There exists $k = k(\mathcal{S})$ such that for any pseudo-Anosov $f \in \mathcal{MCG}(\mathcal{S})$, any vertex $v \in \mathcal{C}(\mathcal{S})$, and any $n \in \mathbb{Z}$,*

$$d_{\mathcal{S}}\big(v, f^n(v)\big) \geq k|n|.$$

The proof of δ-hyperbolicity of $\mathcal{C}(\mathcal{S})$ for a generic $\mathcal{S}$ is nontrivial. We also refer to [Bow06] for an alternate proof.

2.4 Subsurface projections

In this section, we restrict our discussion to domains of an ambient surface $\mathcal{S}$ with $\xi(\mathcal{S}) \geq 1$. Given two domains Y and Z of $\mathcal{S}$, let $Y \cap Z$ represent essential intersections between Y and Z. Intersection with an annular subsurface will be with respect to its core curve. Two domains Y and Z are *disjoint* if $Y \cap Z = \emptyset$; *nested* if $Y \cap Z$ is either Y or Z; and *interlock* if they are neither disjoint or nested. Note that in our definition it is always true that $Y \cap \partial Y = \emptyset$.

Let $Y \subset \mathcal{S}$ be a proper domain. There is a map

$$\pi_Y : \mathcal{C}(\mathcal{S}) \to \mathcal{P}\big(\mathcal{C}(Y)\big),$$

taking an element of $\mathcal{C}(\mathcal{S})$ to a subset of $\mathcal{C}(Y)$ of bounded diameter. We call $\pi_Y(\alpha)$ the *projection* of α to Y. Note that in the definition below, the projection map also makes sense if we replace $\mathcal{S}$ by any subsurface Z which contains Y as a proper subsurface.

We first define the projection to a non-annular domain Y. If $\alpha \cap Y = \emptyset$, then $\pi_Y(\alpha) = \emptyset$. If $\alpha \cap Y = \alpha$, then $\pi_Y(\alpha) = \{\alpha\}$. In all other cases, $\alpha \cap Y$ consist of a collection of arcs in Y. The endpoints of each $\delta \in \alpha \cap Y$ lie on one or two components of ∂Y. Let N be a regular neighborhood of the union of δ with its corresponding component(s) in ∂Y. N has either one or two components, at least one of which is essential in Y. Let $\pi_Y(\delta)$ be the set of essential boundary component(s) of N. We let

$$\pi_Y(\alpha) = \bigcup_{\delta \in \alpha \cap Y} \pi_Y(\delta).$$

Now suppose Y is an annulus with core curve γ. There is a unique cyclic cover of $\mathcal{S}$

$$p : \widehat{Y} \to \mathcal{S}$$

to which Y lifts homeomorphically. Since $\mathcal{S}$ admits a hyperbolic metric, this cover has a natural compactification, also denote by $\widehat{Y}$. We will define $\mathcal{C}(Y) = \mathcal{C}(\widehat{Y})$. For any curve α in $\mathcal{S}$, components of $p^{-1}(\alpha)$ that are essential arcs form a subset in $\mathcal{C}(Y)$. We will let $\pi_Y(\alpha)$ be this corresponding set in $\mathcal{C}(Y)$.

We would like to measure the complexity between curves or curve systems relative to a subsurface. Let $\operatorname{diam}_Y(\cdot)$ be the diameter of subsets in $\mathcal{C}(Y)$. For any two subsets $A, B \subset \mathcal{C}(Y)$, we define

$$d_Y(A, B) = \operatorname{diam}_Y(A \cup B).$$

Given a pair of curves $\alpha, \beta \in \mathcal{C}(\mathcal{S})$, we will define their distance in Y to be

$$d_Y(\alpha, \beta) = d_Y\big(\pi_Y(\alpha), \pi_Y(\beta)\big).$$

For any multicurve σ, one can also project σ to $\mathcal{C}(Y)$ in the obvious way: $\pi_Y(\sigma) = \cup_{\alpha\in\sigma}\pi_Y(\alpha)$. The distance $d_Y(\sigma, \tau)$ is similarly defined. The follow result asserts that subsurface projections are coarsely well-defined and Lipschitz.

Lemma 2.4.1 ([MM00, Lemma 2.3])**.** *For any multicurve σ on $\mathcal{S}$ and any domain $Y \subset \mathcal{S}$, if $\pi_Y(\sigma) \neq \emptyset$, then* $\operatorname{diam}_Y\big(\pi_Y(\sigma)\big) \leq 2$. *Thus, for any pair of simplices σ and β in $\mathcal{C}(\mathcal{S})$,*

$$d_Y(\sigma, \tau) \leq 2d_{\mathcal{S}}(\sigma, \tau).$$

Suppose Y and Z are subsurfaces of $\mathcal{S}$ satisfying $Y \subset Z$. Then the maps π_Y and $\pi_Y \circ \pi_Z$ are coarsely equal as maps from $\mathcal{C}(\mathcal{S}) \to \mathcal{P}(\mathcal{C}(Y))$.

Lemma 2.4.2 ([BKMM06, Lemma 2.12])**.** *There exists a constant M depending only on $\mathcal{S}$ such that for any multicurve σ,*

$$\operatorname{diam}_Y\big(\pi_Y(\sigma), \pi_Y \circ \pi_Z(\sigma)\big) \leq M.$$

Furthermore, we have the following contraction property for the projection map from [MM00, Theorem 3.1].

Theorem 2.4.3 (Bounded geodesic image)**.** *There exists a constant M_0 depending only on $\mathcal{S}$ such that the following holds. Suppose $Y \subset \mathcal{S}$ is a proper essential subsurface, and g is a geodesic in $\mathcal{C}(\mathcal{S})$ such that $\pi_Y(v) \neq \emptyset$ for every vertex $v \in g$. Then*

$$\operatorname{diam}_Y(g) \leq M_0.$$

We say *g cuts Y* if $\pi_Y(v) \neq \emptyset$ for every vertex $v \in g$, and *g misses Y* otherwise. If $d_{\mathcal{S}}(g, \partial Y) \geq 2$ then g cuts Y. On the other hand, by Theorem 2.4.3, if $u, v \in \mathcal{C}(\mathcal{S})$ has $d_Y(u, v) > M_0$, then any geodesic g in $\mathcal{C}(\mathcal{S})$ between u and v misses Y.

2.5 Marking graph

Another useful combinatorial object which admits an action by $\mathcal{MCG}(\mathcal{S})$ is the *marking graph* $\operatorname{Mark}(\mathcal{S})$ of $\mathcal{S}$. Roughly, a *marking* μ on $\mathcal{S}$ is a multicurve c on $\mathcal{S}$ with additionally a set of transverse curves which serve to record twisting data about each curve in c. Below, we give a precise definition that works for any surface $\mathcal{S}$ with $\xi(\mathcal{S}) \geq 1$.

A marking μ on $\mathcal{S}$ is a set of ordered pairs $\{(\alpha_i, t_i)\}$, where the *base curves base*$(\mu) = \{\alpha_i\}$ is a multicurve on $\mathcal{S}$, and each *transversal* t_i is either empty or is a diameter-1 set of vertices in $\mathcal{C}(\alpha_i)$.

The set of transversals $\{t_i\}$ is denoted by $\operatorname{trans}(\mu)$. A transversal t in the pair (α, t) is called *clean* if $t = \pi_\alpha(\beta)$, where β is a curve on $\mathcal{S}$ such that α and β are Farey-neighbors in the subsurface that they fill. A marking μ is *clean* if every non-empty transversal t is clean, and the curve β inducing t does not intersect any other base curve other than α. A marking μ is called *complete* if $\operatorname{base}(\mu)$ is a pants decomposition of $\mathcal{S}$ and no transversal is empty. If μ is complete and clean, then a transversal t determines uniquely the curve β such that $t = \pi_\alpha(\beta)$. If μ is not clean then there is bounded number ways of picking a *compatible* clean marking μ', in the following sense:

Lemma 2.5.1 ([MM00, Lemma 2.4])**.** *There exists a constant M depending only on $\mathcal{S}$ satisfying the following. For any complete marking μ on $\mathcal{S}$. There exists a uniformly bounded number of complete clean markings μ' such that* $\operatorname{base}(\mu) = \operatorname{base}(\mu')$*, and $d_\alpha(t, t') \le M$ for any $(\alpha, t) \in \mu$ and $(\alpha, t') \in \mu'$.*

Definition 2.5.2 (Marking graph)**.** The *marking graph* $\operatorname{Mark}(\mathcal{S})$ is the graph with vertices representing complete clean markings on $\mathcal{S}$. Two vertices $\mu = \{(\alpha_i, \pi_{\alpha_i}(\beta_i))\}$ and $\mu' = \{(\alpha'_i, \pi_{\alpha'}(\beta'_i))\}$ are connected by an edge if they differ by one of the following *elementary moves*:

- *Twist*: For some i, β'_i is obtained from β by a twist or half-twist along α_i. All base curves and other transversals of μ and μ' agree.
- *Flip*: Let μ'' be the (unclean) marking obtained from μ by flipping $(\alpha_i, \pi_{\alpha_i}(\beta_i))$ to $(\beta_i, \pi_{\beta_i}(\alpha_i))$, for some i. Let β_j be the curve intersecting β_i. The marking μ' is a clean marking compatible with μ'' replacing β_j.

We equip $\operatorname{Mark}(\mathcal{S})$ with the combinatorial edge metric, denoted by $d_M = d_{\operatorname{Mark}(\mathcal{S})}$. Like $\mathcal{C}(\mathcal{S})$, $\operatorname{Mark}(\mathcal{S})$ is connected and admits an action of $\mathcal{MCG}(\mathcal{S})$ by isometries. But unlike $\mathcal{C}(\mathcal{S})$, $\operatorname{Mark}(\mathcal{S})$ is locally finite and the action of $\mathcal{MCG}(\mathcal{S})$ is proper. The quotient $\operatorname{Mark}(\mathcal{S})/\mathcal{MCG}(\mathcal{S})$ is also finite, since there are only finitely many complete clean markings up to homeomorphisms of $\mathcal{S}$ [MM00]. Thus, a standard application of Švarc-Milnor implies the orbit map $\mathcal{MCG}(\mathcal{S}) \to \operatorname{Mark}(\mathcal{S})$ is an quasi-isometry.

Definition 2.5.3 (Projection of markings)**.** Let $Y \subset \mathcal{S}$ be essential and let $\mu \in \operatorname{Mark}(\mathcal{S})$. We can project μ to Y, also denoted by $\pi_Y(\mu)$, in the following way. Namely, if Y is not a curve in $\operatorname{base}(\mu)$, then $\pi_Y(\mu) = \pi_Y(\operatorname{base}(\mu))$. If $Y = \alpha \in \operatorname{base}(\mu)$, then we define $\pi_Y(\mu) = t$. Note that, since μ is a complete marking, the projection map is always non-empty. Since $\operatorname{base}(\mu)$ is a diameter-1 set in $\mathcal{C}(\mathcal{S})$, in light of Lemma 2.4.1 the projection map is Lipschitz:

Lemma 2.5.4 ([MM00, Lemma 2.5])**.** *For any $\mu, \nu \in \operatorname{Mark}(\mathcal{S})$ and any domain $Y \subseteq \mathcal{S}$,*

$$d_Y(\mu, \nu) \le 4\, d_M(\mu, \nu).$$

If c is a multicurve and μ a marking such that $c \subseteq \operatorname{base}(\mu)$, then we say μ is an *extension* of c. We will often start with a multicurve c and *extend* it to a marking μ. This amounts to choosing a marking on all the essential non-annular components of $\mathcal{S} \setminus c$ and choosing a transversal for each curve $\alpha \in c$. There are many ways to extend a marking in general, but most often we will need the marking μ to satisfy certain desired properties so those choices will be bounded.

Definition 2.5.5 (Induced Marking)**.** Let $Y \subset \mathcal{S}$ be an non-annular domain. We define a map

$$\Pi_Y : \operatorname{Mark}(\mathcal{S}) \to \operatorname{Mark}(Y).$$

For each marking μ on $\mathcal{S}$, choose a pants decomposition b of Y such that b has minimal intersection with $\pi_Y(\mu)$. We extend b to a marking $\nu = \Pi_Y(\mu)$ on Y as follows. For each curve $\alpha \in b$, choose transversal t_α in Y such that $d_\alpha\big(t_\alpha, \operatorname{trans}(\mu)\big)$ is minimal. The marking $\nu = \{(\alpha, t_\alpha) : \alpha \in b\}$ will be called an *induced marking* of μ on Y, and it is well-defined up to a bounded number of choices. It follows from Lemma 2.4.2 and Lemma 2.5.4 that for any marking μ and any non-annular domain $Y \subset \mathcal{S}$,

$$d_Z\big(\mu, \Pi_Y(\mu)\big) \leq M, \tag{1}$$

where $Z \subset Y$ and M depends only on $\mathcal{S}$.

Definition 2.5.6 (Relative marking extension)**.** Let $\mu \in \mathrm{Mark}(\mathcal{S})$ and c be a multicurve on $\mathcal{S}$. We extend c to a marking $\mu' \in \mathrm{Mark}(\mathcal{S})$ *relative* to μ as follows. For each essential non-annular subsurface $Y \in \mathcal{S} \setminus c$, choose an induced marking $\Pi_Y(\mu)$ on Y. Then for each curve $\alpha \in c$, choose a transversal t_α which has minimal $d_\alpha(t_\alpha, \mathrm{trans}(\mu))$. The union of $\{(\alpha, t_\alpha) : \alpha \in c\}$ with the set of induced markings $\Pi_Y(\mu)$ forms a marking $\mu' \in \mathrm{Mark}(\mathcal{S})$ which is well-defined up to a bounded number of choices.

The following is an immediate consequence of our construction. The point is, except for the possibility that c and $\mathrm{base}(\mu)$ are far in $\mathcal{C}(\mathcal{S})$, μ and μ' look about the same.

Lemma 2.5.7. *Let c be a multicurve on $\mathcal{S}$, μ any marking, and μ' an extension of c relative to μ. For any subsurface $Z \subsetneq \mathcal{S}$, if ∂Z is disjoint from c, or if Z is a curve in c, then*

$$d_Z(\mu', \mu) \leq M,$$

where M depends only on $\mathcal{S}$.

2.6 Hierarchies

In the previous section, we introduced the marking graph $\mathrm{Mark}(\mathcal{S})$ which is quasi-isometric to $\mathcal{MCG}(\mathcal{S})$. In this section, we will introduce the theory of hierarchies, which is useful for constructing efficient paths in $\mathrm{Mark}(\mathcal{S})$. These paths are naturally associated to efficient representations of elements in $\mathcal{MCG}(\mathcal{S})$ in terms of the generators, thus justifying $\mathrm{Mark}(\mathcal{S})$ as a good combinatorial model for $\mathcal{MCG}(\mathcal{S})$.

The idea of hierarchies is to associate to every pair of markings a family of geodesics in curve complexes which behave well with subsurface projections. In order for the theory to work, we need to impose a condition on geodesics in curve complexes called *tightness.* Let $Y \subseteq \mathcal{S}$ be a domain. A tight geodesic g in $\mathcal{C}(Y)$ is a sequence $\{v_0, \ldots, v_n\}$ of simplices in $\mathcal{C}(Y)$, such that any sequence of vertices in g is a geodesic in $\mathcal{C}(Y)$ in the usual sense, and $v_{i-1} \cup v_{i+1}$ fill a subsurface $Z \subset Y$ such that $\partial Z = v_i$. We remark that the original definition [MM00, Definition 4.2] consists of more information. It is a theorem of Masur-Minsky that any two points in $\mathcal{C}(Y)$ is connected by at least one and at most finitely many tight geodesics [MM00, Lemma 4.5 and Corollary 6.14]. Henceforth, a geodesic in a curve complex will always mean a tight geodesic. By an abuse of notation, we will refer to v_i's as vertices of g. We will say the *length* of g is n, and write $|g| = n$. We will say Y is the *domain* or *support* of g, and write $D(g) = Y$. We will sometimes use the notation $[v_0, v_n]$ to mean any geodesic from v_0 to v_n in $\mathcal{C}(Y)$. Since $\mathcal{C}(Y)$ is δ-hyperbolic, all (finitely many) geodesics from v_0 to v_n are fellow-travelers.

A *hierarchy* on $\mathcal{S}$ is a collection H of geodesics such that each geodesic $g \in H$ is supported on some domain $Y \subset \mathcal{S}$, with a distinguished *main geodesic* $g_H = [v_0, v_n]$ supported on $\mathcal{S}$. Additionally, H comes equipped with a pair of markings $I(H)$ and $T(H)$ on $\mathcal{S}$, called the *initial marking* and the *terminal marking* of H, respectively, such that $v_0 \subseteq \mathrm{base}(I(H))$ and $v_n \subseteq \mathrm{base}(T(H))$. The *length* of H is $|H| = \sum_{h \in H} |h|$. We will usually assume $I(H)$ and $T(H)$ are complete clean marking on $\mathcal{S}$. To be a hierarchy, the geodesics in H must satisfy certain technical conditions called *subordinacy.* Roughly, given a geodesic g in $\mathcal{C}(\mathcal{S})$, one can inductively construct a hierarchy H with $g = g_H$. For each vertex v_i in g_H, the vertices v_{i-1} to v_{i+1} are contained in some component Z of $\mathcal{S} \setminus v_i$. The geodesic $h = [v_{i-1}, v_{i+1}]$ in $\mathcal{C}(Z)$ will be an element of H and is subordinate to

g_H. One can continue with this process with all vertices of g_H and with h and so on. See [MM00, Definition 4.4] for a precise definition of hierarchies. We list some properties of hierarchies below, after the following definition.

Definition 2.6.1. Given a non-annular domain $Y \subseteq \mathcal{S}$, and a simplex $v \in \mathcal{C}(Y)$, we say Z is a *component domain* of (Y, v) if Z is a component of $Y \setminus v$ which is not a pair of pants or if Z is a curve in v.

Theorem 2.6.2. *The following statements hold for hierarchies.*

1 *(Existence) Given any markings μ and ν on $\mathcal{S}$, there exists a hierarchy H with $I(H) = \mu$ and $T(H) = \nu$ [MM00, Theorem 4.6].*

2 *(Uniqueness of geodesics) For any hierarchy H, if $h, h' \in H$ have $D(h) = D(h')$, then $h = h'$ [MM00, Theorem 4.7].*

3 *(Completeness) For every pair, $h \in H$ and a vertex $v \in h$, if Y is a component domain of $(D(h), v)$, then Y is domain for a geodesic $k \in H$ [MM00, Theorem 4.20].*

We will sometimes denote an hierarchy from μ to ν by $H(\mu, \nu)$. The following lemma explains the relationship between a geodesic $h \in H$ and the projection of $I(H)$ and $T(H)$ to $D(h)$.

Lemma 2.6.3 ([MM00, Lemma 6.2])**.** *There exist constants $M_1 > M_2$, depending only on $\mathcal{S}$, such that if H is any hierarchy in $\mathcal{S}$ and*

$$d_Y\big(I(H), T(H)\big) \geq M_2$$

for a subsurface Y in $\mathcal{S}$, then Y is a domain for a geodesic $h \in H$.

Conversely, if $h \in H$ is any geodesic with $Y = D(h)$, then $d_Y\big(I(H), T(H)\big)$ and h are fellow travelers with a uniform constant. In particular,

$$\big||h| - d_Y\big(I(H), T(H)\big)\big| \leq M_1.$$

For any pair of markings $\mu, \nu \in \mathrm{Mark}(\mathcal{S})$, we will call a domain Y a *large link* for μ and ν if $d_Y(\mu, \nu) \geq M_2$.

The following theorem summarizes two results that are vital to this paper. To simplify the statements we introduce some notations which we will adopt for rest of the paper. Below, $\mathfrak{a}$ and $\mathfrak{b}$ represent quantities such as distances or lengths, and k and c are constants that depend only on $\mathcal{S}$ (unless otherwise noted).

Notations 2.6.4. 1. If $\mathfrak{a} \leq k\mathfrak{b} + c$, we say a is *coarsely bounded* by b, and write $\mathfrak{a} \prec \mathfrak{b}$

2. If $\frac{1}{k}\mathfrak{b} - C \leq \mathfrak{a} \leq k\mathfrak{b} + c$, we say a is *coarsely equal* to b, and write $\mathfrak{a} \asymp \mathfrak{b}$.

In the following, the coarse equality on the left is [MM00, Theorem 6.10]. The coarse equality on the right is called the distance formula [MM00, Theorem 6.12].

Theorem 2.6.5. *There exists a constant L_0 depending only on S such that, for any $L \geq L_0$ and any $\mu, \nu \in \mathrm{Mark}(\mathcal{S})$ and any hierarchy $H = H(\mu, \nu)$,*

$$|H| \quad \asymp \quad d_M(\mu, \nu) \quad \asymp \sum_{\substack{Y \subseteq \mathcal{S} \\ d_Y(\mu,\nu) \geq L}} d_Y(\mu, \nu).$$

On the right, the constants involved in $\asymp$ depend on L.

Fix a generating set Λ for $\mathcal{MCG}(\mathcal{S})$. To realize the quasi-isometry between $\mathrm{Mark}(\mathcal{S})$ and $\mathcal{MCG}(\mathcal{S})$, we fix a base marking μ_0 in $\mathrm{Mark}(\mathcal{S})$. Then $d_M(\mu_0, f\mu_0) \asymp |f|$, with constants depending only on μ_0 and Λ. The following is an immediate consequence of Theorem 2.6.5.

Corollary 2.6.6 ([MM00, Theorem 7.1])**.** *Let* $\mu_0 \in \mathrm{Mark}(\mathcal{S})$ *be a fixed base marking. For any element* $f \in \mathcal{MCG}(\mathcal{S})$*,*

$$|H(\mu_0, f\mu_0)| \quad \asymp \quad |f|.$$

Let $Y \subset \mathcal{S}$ be a proper non-annular domain. There is a coarse embedding $\mathrm{Mark}(Y) \xrightarrow{j} \mathrm{Mark}(\mathcal{S})$: fix a marking in each complementary component of $\mathcal{S} \setminus Y$ and a transversal to each curve in ∂Y. The map j sends the marking $\nu \in \mathrm{Mark}(Y)$ in an obvious way such that $\partial Y \subseteq \mathrm{base}\big(j(\nu)\big)$ and for all $\nu_1, \nu_2 \in \mathrm{Mark}(Y)$,

$$d_{\mathrm{Mark}(Y)}(\nu_1, \nu_2) \asymp d_M\big(j(\nu_1), j(\nu_2)\big). \tag{2}$$

Equation (2) follows from the distance formula and one can make the coarse constants independent of Y and j.

2.7 Slices

The connection between paths in $\mathrm{Mark}(\mathcal{S})$ and hierarchies come from *slices* of a hierarchy. The following definition comes from from [MM00, §5].

Definition 2.7.1 (Slices)**.** A *(complete) slice* of a hierarchy H is a set τ of *pointed geodesics* (h, v) in H, i.e. $h \in H$ and v is a vertex of h, satisfying the following properties:

(S1) Any geodesic h of H appears at most once in τ.

(S2) There is a distinguished pair, the *bottom pair*, (g_H, b) of τ.

(S3) For every $(k, w) \in \tau$ other than the bottom pair, $D(k)$ is a component domain of $(D(h), v)$ for some $(h, v) \in \tau$.

(S4) Given $(h, v) \in \tau$, for every component domain Y of $(D(h), v)$ there is a pair $(k, w) \in \tau$ with $D(k) = Y$.

The *initial slice* τ_0 of H is one where every pair $(h, v) \in \tau$ has v the first vertex of h. In particular, the main geodesic g_H and its initial vertex is a pair in τ_0, and τ_0 can be constructed inductively using the axioms of slices. Similarly, the *terminal slice* of H is defined. To any slice τ we can associate a complete marking μ_τ as follows. First let μ be the marking with

$$\mathrm{base}(\mu) = \{\ v \ : \ (h, v) \in \tau \text{ and } D(h) \text{ is not an annulus }\}.$$

For each base curve α, if $(k, t) \in \tau$ is such that k is a geodesic in $\mathcal{C}(\alpha)$, then let t be the transversal to α in μ. The marking μ is complete but not necessarily clean. Any clean marking μ_τ compatible with μ will be called a *compatible* marking with τ. By Lemma 2.5.1, the number of choices for μ_τ is bounded. $I(H)$ and $T(H)$ are respectively compatible markings with the initial and terminal slice of H.

Given any slice τ in H, there is a notion of *(forward) elementary move* on τ which is roughly moving a vertex v of some pair $(h, v) \in \tau$ forward by one step in the geodesic h to obtain a new slice τ'. We write $\tau \to \tau'$. See [MM00, §5] for a precise definition. If μ and μ' are compatible marking with τ and τ', then by [MM00, Lemma 5.5], $d_M(\mu, \mu') \prec 1$. We will write $\mu \to \mu'$ to mean any path in $\mathrm{Mark}(\mathcal{S})$ connecting μ to μ'. To prove$|H| \asymp d_M(I(H), T(H))$, Masur and Minsky in

[MM00] established the existence of a *resolution* of H, which is a sequence of forward elementary moves

$$\tau_0 \to \cdots \to \tau_n,$$

where τ_0 is the initial slice and τ_n is the terminal slice of H. For each τ_i in the resolution, let μ_i be a compatible marking with τ_i. The corresponding path in $\mathrm{Mark}(\mathcal{S})$

$$I(H) = \mu_0 \to \cdots \to \mu_n = T(H)$$

turns out to be a quasi-geodesic with uniform constants, and $d_M\big(I(H), T(H)\big) \asymp n \asymp |H|$. An important fact in [Min02] is that, for any slice τ in H, there is a resolution of H containing τ. We will call any marking compatible to some slice in a hierarchy H a *hierarchal marking.* The following statements are true for hierarchal markings.

Lemma 2.7.2. *There exists a constant M depending only on $\mathcal{S}$ such that the following hold. Let H be a hierarchy. If $\mu \in \mathrm{Mark}(\mathcal{S})$ is a hierarchal marking for H, then*

$$d_M\big(I(H), \mu\big) + d_M\big(\mu, T(H)\big) \leq d_M\big(I(H), T(H)\big) + M$$

Furthermore, for any $Y \subseteq \mathcal{S}$,

$$d_Y\big(I(H), \mu\big) + d_Y\big(\mu, T(H)\big) \leq d_Y\big(I(H), T(H)\big) + M$$

2.8 Time order

The geodesics or domains of geodesics in a hierarchy H satisfy a partial order $<_t$, called *time order.* We refer to [MM00, §4.6] for the definition. The idea comes from the observation that the vertices of a geodesic g are linearly ordered: $v_i < v_j$ if $i < j$. Combining this observation with the subordinacy structure on H, one can try to order a pair of geodesics $h, h' \in H$. A simple situation is if $D(h)$ is a component domain of $(D(g), v_i)$ and $D(h')$ is a component domain of $(D(g), v_j)$, then $D(h) <_t D(h')$. If $D(h)$ and $D(h')$ are different components of $(D(g), v_i)$, then in fact they are not time-ordered. The situation gets more complicated if h and h' are not subordinate to the same geodesic g. An important fact is that if $D(h)$ and $D(h')$ interlock, then they must be time-ordered. If $D(h)$ and $D(h')$ are nested, then they cannot be time-ordered. The ambiguous case is when $D(h)$ and $D(h')$ are disjoint; sometimes they are time-ordered and sometimes not. The issue of disjoint domains will come up in this paper. We summarize the above results and state some useful consequences of time order.

Theorem 2.8.1 ([MM00, Lemma 4.18 and 4.19])**.** *There exists a relation $<_t$, called* time-order*, on domains of geodesics in H such that:*

- *The relation $<_t$ is a strict partial order.*
- *If h and h' are geodesics in H such that $Y = D(h)$ and $Z = D(h')$ interlock, then either $Y <_t Z$ or $Z <_t Y$.*
- *If $Y \subset Z$, then Y and Z are not time-ordered.*
- *If Y and Z are such that they lie in different component domains of $(D(m), v)$, for some geodesic m in H, then Y and Z are not time-ordered.*

The constant M_1 of Lemma 2.6.3 can be chosen so that following hold.

Lemma 2.8.2 ([MM00, Lemma 6.11])**.** *Let H be a hierarchy. Suppose Y and Z are domains for geodesics in H such that Y and Z interlock. If $Y <_t Z$, then $d_Y\big(\partial Z, T(H)\big) \leq M_1$ and $d_Z\big(I(H), \partial Y\big) \leq M_1$.*

Using slices, the constant M_1 can be chosen so the following version of Lemma 2.8.2 also holds.

Lemma 2.8.3. *With the same hypothesis as above. There exists a hierarchy marking* ν *such that* $d_Y\big(\nu, T(H)\big) \leq M_1$ *and* $d_Z(I(H), \nu) \leq M_1$*.*

We also have:

Corollary 2.8.4 ([BM08, Lemma 1])**.** *With the same hypothesis as above. For any marking* $\mu \in \mathrm{Mark}(\mathcal{S})$*, either* $d_Y\big(\mu, T(H)\big) \leq 2M_1$ *or* $d_Z(I(H), \mu) \leq 2M_1$*.*

2.9 Separating Marking

The following definition and lemma do not explicitly appear in [MM00]. Although the lemma is a direct consequence of hierarchies, we offer a brief sketch of its proof.

Definition 2.9.1 (Separating marking)**.** Let H be a hierarchy. A slice τ is called a *separating slice* if for every pair $(h, v) \in \tau$, with $h \neq g_H$, has the property that v is the terminal vertex of h. We remark that once the bottom pair (g_H, b) is fixed, then the separating slice τ containing (g_H, b) is uniquely determined by the axioms of slices. In particular, if b is the terminal vertex of g_H, then τ is the terminal slice of H. If τ is a separating slice containing (g_H, b), then any marking μ compatible with τ is called a *separating marking* at b.

Lemma 2.9.2. *The constant* M_1 *of Lemma 2.6.3 can be chosen so that following hold. Let* H *be a hierarchy. Let* b *be any vertex in* g_H *and let* μ *be a separating marking at* b*. Then for any proper domain* $Y \subset \mathcal{S}$*, either* $d_Y(I(H), \mu) \leq M_1$ *or* $d_Y(\mu, T(H)) \leq M_1$*.*

Proof. We may assume $Y \subset \mathcal{S}$ has $d_Y\big(I(H), T(H)\big) > M_1$. Since $M_1 \geq M_2$, Y is a domain for a geodesic $h_Y \in H$. Without a loss of generality, we may assume Y is a component domain of (g_H, c). If c appears before b along g_H, then $d_Y\big(\mu, T(H)\big) \leq M_1$. Similarly, if c appears after b along g_H, then $d_Y(I(H), \mu) \leq M_1$. Both of these facts can be seen as a consequence of Lemma 2.8.2. The remaining case is $b = c$. In this case, τ must contain (h_Y, v), where v is the terminal vertex of h_Y. Therefore, it must be that $d_Y\big(\mu, T(H)\big) \leq M_1$. □

Remark 2.9.3. In our definition of separating slice, the preference for terminal vertices is arbitrary. Lemma 2.9.2 would remain true if we allowed only initial vertices or a mixture of initial and terminal.

2.10 Collecting constants

For the rest of the paper, we will fix the following set of constants. Let N be the constant of Lemma 3.1.1. Let M_0 be the constant of Theorem 2.4.3. Let L_0 be the constant of Theorem 2.6.5. Let M_1 and M_2 be the constants coming from Lemma 2.6.3. We will also fix one constant M_3 for Lemma 2.4.2, Lemma 2.5.1, Equation (1), Lemma 2.5.7, and Lemma 2.7.2. We may assume $M_1 \geq M_2, M_3$. In addition, since up to homeomorphism there are only finitely many subsurfaces of $\mathcal{S}$, we can choose a hyperbolicity constant δ which works for all $\mathcal{C}(Z)$, $Z \subseteq \mathcal{S}$.

3 Two technical lemmas

This section contains some technical results about finite order mapping classes.

To prove L.B.C. property, we need to understand the geometry of the action of finite order mapping classes on $\mathrm{Mark}(\mathcal{S})$. The first observation is that finite order elements act on $\mathrm{Mark}(\mathcal{S})$ with coarse fixed points. We will eventually prove that the action has the property that the translation distance of a finite order element f, or $d_M(\mu_0, f\mu_0)$ where μ_0 is the base marking, is coarsely bounded by the distance from μ_0 to the fixed point sets of f. In other words, finite order elements of $\mathcal{MCG}(\mathcal{S})$ act *elliptically* on $\mathrm{Mark}(\mathcal{S})$.

In this section, we consider what happens if a fixed point μ of f is far from μ_0 relative to the translation distance of f. By the distance formula, there must be some $X \subseteq S$ such that $d_X(\mu_0, \mu)$ is large relative to $d_X(\mu_0, f\mu_0)$. With some additional conditions, X will be called a *bad domain* for μ and we will prove a structure theorem for the set of bad domains in a hierarchy $H(\mu_0, \mu)$. In this next section, we will use this structure theorem to construct a coarse fixed point of f close to μ_0 relative to the translation distance of f. From there, we can derive L.B.C. property for finite order mapping classes by a standard argument.

3.1 Fixed points and symmetric points

We state some useful facts about finite order mapping classes below.

Lemma 3.1.1. *There exists a constant N, depending only on $\mathcal{S}$, such any finite order element $f \in \mathcal{MCG}(\mathcal{S})$ has* $\mathrm{order}(f) \leq N$.

Lemma 3.1.2. *There are finitely many conjugacy classes of finite order elements in $\mathcal{MCG}(\mathcal{S})$.*

Definition 3.1.3. Let $f \in \mathcal{MCG}(\mathcal{S})$ be of finite order. We define the set of *r-fixed points* of f as

$$\mathrm{Fix}_r(f) = \{ \ \mu \in \mathrm{Mark}(\mathcal{S}) \ \ : \ \ d_M(\mu, f\mu) \leq r \ \}.$$

Also, define the set of *r-symmetric points* for f to be

$$\widetilde{\mathrm{Fix}}_r(f) = \{ \ \mu \in \mathrm{Mark}(\mathcal{S}) \ \ : \ \ d_Y(\mu, f\mu) \leq r, \forall Y \subseteq \mathcal{S} \ \}.$$

Lemma 3.1.4. *There exist constants R_1, depending only on $\mathcal{S}$, such that* $\mathrm{Fix}_{R_1}(f) \neq \emptyset$ *and* $\widetilde{\mathrm{Fix}}_{R_1}(f) \neq \emptyset$, *for any finite order element $f \in \mathcal{MCG}(\mathcal{S})$.*

Proof. We first show $\mathrm{Fix}_{R_1}(f) \neq \emptyset$. By Nielsen's Realization [Ker83], f fixes a hyperbolic metric ρ on $\mathcal{S}$. Let $\mu \in \mathrm{Mark}(\mathcal{S})$ be a marking such that the sum of the hyperbolic lengths of curves in μ measured in ρ is minimal over $\mathrm{Mark}(\mathcal{S})$. Let $R_f = d_M(\mu, f\mu)$. If $g = \omega f \omega^{-1}$ for some $\omega \in \mathcal{MCG}(\mathcal{S})$, then g fixes the metric $\omega_* \rho$. Hence $R_g = R_f$.

Now, using Lemma 3.1.2, we can let R_1 be the maximum of the constants R_f ranging over all conjugacy classes. By the distance formula, one can choose R_1 such that $\widetilde{\mathrm{Fix}}_{R_1}(f) \neq \emptyset$ as well. $\square$

Remark 3.1.5. The set $\mathrm{Fix}_R(f)$ is a quasi-convex subset of $\mathrm{Mark}(\mathcal{S})$. That is, there exists a constant $\epsilon = \epsilon(R)$, such that for any $\mu, \nu \in \mathrm{Fix}_R(f)$, any geodesic $[\mu, \nu]$ in $\mathrm{Mark}(\mathcal{S})$ must stay within an ϵ-neighborhood of $\mathrm{Fix}_R(f)$. In fact, a stronger statement about the geometry of $\mathrm{Fix}_R(f)$ is known. Suppose f fixes the metric ρ. Let X be $\mathcal{S}$ equipped with ρ. The action of f on X is an isometry, so its quotient $\bar{X} = X/f$ is an orbifold. One can coarsely identify $\mathrm{Fix}_R(f)$ with $\mathrm{Mark}(\bar{X})$. The map $X \to \bar{X}$ is a (branched) covering map. By [RS07], the lifting of $\mathrm{Mark}(\bar{X})$ to $\mathrm{Mark}(X) = \mathrm{Mark}(\mathcal{S})$ is a quasi-isometric embedding.

3.2 An example

Before proceeding to the first technical lemma, let's discuss a motivating example. The following refers to Figure 1.

Consider the closed surface $\mathcal{S}_2$ of genus two. Let f be the mapping class of order two which permutes the two holes of S_2. The curves in Figure 1 represent a marking μ on S_2: the red curves are the base curves of μ and the blue curves are the transversal curves of μ. The marking μ is a fixed point of f (in fact, μ is a 0–fixed point of f). Let X and Y be the pair of once-punctured tori in S_2 permuted by f. Since X is once-punctured torus, $\mathcal{C}(X)$ is homeomorphic to the Farey graph $\mathcal{F}$. Markings on X correspond to edges in $\mathcal{F}$, thus we may represent the projection $\nu_1 = \pi_Y(\mu)$ by an edge in $\mathcal{F}$. After identifying $\mathcal{C}(X)$ and $\mathcal{C}(Y)$ via the map f, the projection $\nu_2 = \pi_Y(\mu)$ is represented by the same edge in $\mathcal{F}$.

We can construct some other coarse fixed points of f as follows. Choose any marking ν on X. Let α be the separating curve bounding X and Y. Let $t = \pi_\alpha(\mu)$ be the transversal to α associated to μ. Any clean marking μ' compatible with $\nu \cup f(\nu) \cup \alpha \cup t$ has the property that $d_M(\mu', f\mu') \prec 1$.



Figure 1: On the left, the curves represent a fixed marking μ for the order two element $f \in \mathcal{MCG}(\mathcal{S}_2)$ that permutes the holes of $\mathcal{S}_2$. On the right is the Farey graph $\mathcal{F}$ which is isomorphic to $\mathcal{C}(X)$ and $\mathcal{C}(Y)$. Markings on X or Y correspond to edges in $\mathcal{F}$. Here, $\nu_1 = \pi_X(\mu)$ and $\nu_2 = \pi_Y(\mu)$, and $z_1 = \pi_X(\mu_0)$ and $z_2 = \pi_Y(\mu_0)$, where μ_0 is the fixed based marking.

Now suppose the base marking μ_0 has the following form. Let z_1 and z_2 be any two edges in $\mathcal{F}$. Suppose μ_0 contains α as a base curve, $d_\alpha(\mu_0) = t$, $d_X(\mu_0) = z_1$ and $d_Y(\mu_0) = z_2$. We will also assume that ν_1 and z_j, $j = 1, 2$ do no have large projections to any annular domain. These assumptions imply that X and Y are the only possible large links for μ_0 and μ. More precisely,

$$d_{\mathrm{Mark}(\mathcal{S})}(\mu_0, \mu) \asymp d_X(\mu_0, \mu) + d_Y(\mu_0, \mu) = d_{\mathcal{F}}(\nu_1, z_1) + d_{\mathcal{F}}(\nu_2, z_2).$$

We want to understand which fixed point of f is close to μ_0. If we are in the situation that

$$d_{\mathcal{F}}(\nu_1, z_1) \le d_{\mathcal{F}}(z_1, z_2), \tag{3}$$

then we also have

$$d_{\mathcal{F}}(\nu_2, z_2) = d_{\mathcal{F}}(\nu_1, z_2) \le d_{\mathcal{F}}(\nu_1, z_1) + d_{\mathcal{F}}(z_1, z_2) \le 2d_{\mathcal{F}}(z_1, z_2) \tag{4}$$

Note that $d_{\mathcal{F}}(z_1, z_2) = d_X(\mu_0, f\mu_0) = d_Y(\mu_0, f\mu_0)$. Hence, μ is already close to μ_0 in the sense that

$$d_M(\mu_0, \mu) \prec d_M(\mu_0, f\mu_0).$$

If, on the other hand,

$$d_{\mathcal{F}}(\nu_1, z_1) > d_{\mathcal{F}}(z_1, z_2) \tag{5}$$

is sufficiently large, then we also have

$$d_{\mathcal{F}}(\nu_2, z_2) \ge d_{\mathcal{F}}(\nu_1, z_1) - d_{\mathcal{F}}(z_1, z_2) \succ d_{\mathcal{F}}(z_1, z_2). \tag{6}$$

In this situation, X is a *bad domain* for μ. The existence of X disqualifies μ from being adequately close to μ_0 in X and Y. We sketch a procedure on how to construct a closer fixed point of f. Start by considering the geodesic path from ν_1 to z_1 in $\mathcal{F}$. Choose a vertex v along the path satisfying $d_{\mathcal{F}}(v, \nu_1) \leq d_{\mathcal{F}}(\nu_1, \nu_2)$. Extend the vertex v to an edge ν in $\mathcal{F}$, with care taken to ensure ν and z_j do not have large projections to annular domains (this requires an inductive procedure). This edge ν serves as the marking on X. Then, as in the previous paragraph, the marking μ' compatible with $\nu \cup f(\nu) \cup \alpha \cup t$ is a coarse fixed point of f. Using the same computations as in (3) and (4), our choice of ν ensures that

$$d_{\mathrm{Mark}(\mathcal{S})}(\mu_0, \mu) \prec d_{\mathrm{Mark}(\mathcal{S})}(\mu_0, f\mu_0).$$

A couple of observations about the example. First note that, for (5) to imply (6), the sufficiently large condition can be made to depend only on the order of f. Secondly, the procedure for constructing the new fixed point μ' affects both X and Y at the same time. The main reason is that X and Y are disjoint. A modification of μ in X is transferable to Y via f. This ensures maximal symmetry for the action f.

In the following and in the subsequent section, we generalize this example. Equations (5) and (6) are motivations behind the definition of a bad domain (see Definition 3.3.1). The structure result for bad domains, Lemma 3.3.4, shows that if X is a bad domain for μ, then X and its orbits are all disjoint and are all large links for μ_0 and μ. In the next section, we will show how to use the disjointness result to cook up the desired fixed point of f close to μ_0, using a procedure similar to the one described above.

3.3 Bad domains and the first technical lemma

We remark that Lemma 3.1.2 does not a priori help us with L.B.C. property as each conjugacy class has infinitely many elements, but it will play an essential role later. In the following, let $\mu_0 \in \mathrm{Mark}(\mathcal{S})$ be the fixed base marking. We recall notations of §2.10 and let R_1 be the minimal constant satisfying Lemma 3.1.4. Set

$$\Theta = 6M_1 + 4\delta. \tag{7}$$

Definition 3.3.1 (Bad domains)**.** For any $X \subset \mathcal{S}$ a proper subsurface, set L_X as the integer such that f^{L_X+1} is the first return map of f to X. Note that for any X, $L_X < \mathrm{order}(f) \leq N$, where N is the constant of 3.1.1. Let $f \in \mathcal{MCG}(\mathcal{S})$ be of finite order. Let $R \geq R_1$ and let $\mu \in \mathrm{Mark}(\mathcal{S})$ be any marking. We say $X \subseteq \mathcal{S}$ is a *R-bad domain* for μ (and f) if

- For $X = \mathcal{S}$,
$$d_{\mathcal{S}}(\mu_0, \mu) > d_{\mathcal{S}}(\mu_0, f\mu_0) + R. \tag{8}$$
- For $X \neq \mathcal{S}$,
$$d_X(\mu_0, \mu) > 2N\Big(\max_{0 \leq i \leq L_X} \big\{d_{f^i(X)}(\mu_0, f\mu_0)\big\}\Big) + NR + \Theta. \tag{9}$$

Denote by $\Omega(\mu, R, f)$ or $\Omega(\mu, R)$ the set of all R-bad domains for μ (and f). Note that if $R' > R$, then $\Omega(\mu, R') \subseteq \Omega(\mu, R)$. We remark that the constant "$2N$" in the definition of bad domains will not play a role until the next section.

Definition 3.3.2 (Partial order on $\Omega(\mu, R)$)**.** We endow $\Omega(\mu, R)$ with a partial order "$\lhd$" following these rules. Let $X, Y \in \Omega(\mu, R)$, and let $H = H(\mu_0, \mu)$ be a fixed hierarchy.

(O1) If $\xi(X) < \xi(Y)$, then $X \lhd Y$. In particular, if $\mathcal{S} \in \Omega(\mu, R)$, then $\mathcal{S}$ is the maximal element.

(O2) If $\xi(X) = \xi(Y)$ and $X <_t Y$ in $H(\mu_0, \mu)$, then $Y \lhd X$.

Definition 3.3.3 (Complexity of $\Omega(\mu, R)$)**.** If $\Omega(\mu, R)$ is nonempty, then the complexity of the maximal element in $\Omega(\mu, R)$ is called the *complexity of* $\Omega(\mu, R)$, denoted by $\xi(\mu, R)$. The minimal complexity over all subsurfaces of $\mathcal{S}$ is -1, coming from an annulus. For convenience, we will let $\xi(\emptyset) = -2$. We make the trivial observations that whenever

$$\Omega(\mu', R') \subseteq \Omega(\mu, R) \qquad \Longrightarrow \qquad \xi(\mu', R') \leq \xi(\mu, R).$$

The following is a consequence of the definition of R-bad domains for μ and f, if μ happens to be a R-symmetric point for f.

Lemma 3.3.4 (Structure of bad domains)**.** *Let $R \geq R_1$ and let $\mu \in \widetilde{\mathrm{Fix}}_R(f)$. If $X \in \Omega(\mu, R)$ and $X \neq \mathcal{S}$, then*

$$X, f(X), \dots f^{L_X}(X)$$

are all domains for geodesics in $H(\mu_0, \mu)$ and are all mutually disjoint.

Proof. Note that $f^{-i}(X) = f^{L_X+1-i}(X)$. We will proceed by induction on $f^{-i}(X)$, for $i = 0, \dots, L_X$. Our assumption is that $X \in \Omega(\mu, R)$ and $X \neq \mathcal{S}$. In particular, this means that $d_X(\mu_0, \mu) > \Theta > M_2$, so X is a domain for a geodesic in $H(\mu_0, \mu)$. This concludes the base case. Let's now assume

$$X, f^{-1}(X), \dots, f^{-n+1}(X)$$

are all domains for geodesics in $H(\mu_0, \mu)$ and are all mutually disjoint. We will show $f^{-n}(X)$ supports a geodesic in $H(\mu_0, \mu)$. Recursively, we have

$$\begin{aligned}
&d_{f^{-n}(X)}(\mu_0, \mu) \\
&\qquad = d_{f^{-n+1}(X)}(f\mu_0, f\mu) \\
&\qquad \geq d_{f^{-n+1}(X)}(\mu_0, \mu) - d_{f^{-n+1}(X)}(\mu_0, f\mu_0) - d_{f^{-n+1}(X)}(\mu, f\mu) \\
&\qquad > d_{f^{-n+2}(X)}(f\mu_0, f\mu) - d_{f^{-n+1}(X)}(\mu_0, f\mu_0) - R \\
&\qquad\qquad \vdots \\
&\qquad \geq d_X(\mu_0, \mu) - \left(\sum_{i=1}^{n} d_{f^{-n+i}(X)}(\mu_0, f\mu_0) \right) - nR \\
&\qquad \geq d_X(\mu_0, \mu) - n\Big(\max_{1 \leq i \leq n} \left\{ d_{f^{-n+i}(X)}(\mu_0, f\mu_0) \right\} \Big) - nR \\
\text{By (9)}\quad &\qquad > (N - n)\Big(\max_{1 \leq i \leq L_X + 1} \left\{ d_{f^{-n+i}(X)}(\mu_0, f\mu_0) \right\} \Big) + (N - n)R + \Theta
\end{aligned}$$

Since $N > L_X \geq n$, we have in particular

$$d_{f^{-n}(X)}(\mu_0, \mu) > \Theta > M_2.$$

Therefore, $f^{-n}(X)$ supports a geodesic in $H(\mu_0, \mu)$.

Now let's prove $f^{-n}(X)$ is disjoint from each $X, \dots, f^{-n+1}(X)$. Observe that $f^{-n}(X)$ and $f^{-i}(X)$ are disjoint if and only if $f^{-n+i}(X)$ and X are disjoint. Hence it is enough to show $f^{-n}(X)$ and X are disjoint. By way of contradiction, let's assume X and $f^{-n}(X)$ are not disjoint. The two domains have the same complexity so they must interlock. They both support geodesics in $H(\mu_0, \mu)$ so, by Theorem 2.8.1, they are time-ordered. We have two cases.

The first case is $X <_t f^{-n}(X)$. As in Lemma 2.8.3, we may choose a hierarchal marking ν in $H(\mu_0, \mu)$ such that

$$d_X(\nu, \mu) \leq M_1 \quad \text{and} \quad d_{f^{-n}(X)}(\mu_0, \nu) \leq M_1, \tag{10}$$

By the triangle inequality,

$$\begin{aligned} d_X(\mu_0, f^n\nu) &\le d_X(\mu_0, f^n\mu_0) + d_X(f^n\mu_0, f^n\nu) \\ &\le \left(\sum_{j=0}^{n-1} d_X(f^j\mu_0, f^{j+1}\mu_0)\right) + d_{f^{-n}(X)}(\mu_0, \nu) \\ &\le \left(\sum_{j=0}^{n-1} d_{f^{-j}(X)}(\mu_0, f\mu_0)\right) + d_{f^{-n}(X)}(\mu_0, \nu) \\ &\le n\left(\max_{0\le j\le n-1} \left\{ d_{f^{-j}(X)}(\mu_0, f\mu_0) \right\}\right) + M_1. \end{aligned} \tag{11}$$

Using the triangle inequality again, along with (9) and (11), we have

$$d_X(f^n\nu, \mu) \ge d_X(\mu_0, \mu) - d_X(\mu_0, f^n\nu) > 2M_1.$$

Therefore, by Corollary 2.8.4,

$$d_{f^{-n}(X)}(\mu_0, f^n\nu) \le 2M_1.$$

Now we consider $d_X(\mu_0, f^{in}\mu)$. By iterating the argument we obtain, for every $i \ge 0$,

$$\begin{aligned} d_X(\mu_0, f^{(i+1)n}\nu) &\le d_X(\mu_0, f^n\mu_0) + d_X(f^n\mu_0, f^{(i+1)n}\nu) \\ &\le n\Big(\max_{0\le j\le n-1} \left\{ d_{f^{-j}(X)}(\mu_0, f\mu_0) \right\}\Big) + d_{f^{-n}(X)}(\mu_0, f^{in}\nu) \\ &\le n\Big(\max_{0\le j\le n-1} \left\{ d_{f^{-j}(X)}(\mu_0, f\mu_0) \right\}\Big) + 2M_1. \end{aligned}$$

Since there is some i for which f^{in} is the identity map, the latter inequality must also hold for $d_X(\mu_0, \nu)$. With this fact coupled with the first half of (10), we derive the following violation to $X \in \Omega(\mu, R)$:

$$\begin{aligned} d_X(\mu_0, \mu) &\le d_X(\mu_0, \nu) + d_X(\nu, \mu) \\ &\le n\Big(\max_{0\le j\le L_X} \left\{d_{f^{-j}(X)}(\mu_0, f\mu_0)\right\}\Big) + 2M_1 + d_X(\nu, \mu) \\ &\le N\Big(\max_{0\le j\le L_X} \left\{d_{f^{-j}(X)}(\mu_0, f\mu_0)\right\}\Big) + 3M_1 \end{aligned}$$

Thus, it is not possible for $X <_t f^{-n}(X)$.

To eliminate the second case $f^{-n}(X) <_t X$, we argue similarly. Now choose a marking ν such that

$$d_X(\mu_0, \nu) \le M_1 \quad \text{and} \quad d_{f^{-n}(X)}(\nu, \mu) \le M_1. \tag{12}$$

Then, using the fact that $\mu \in \widetilde{\mathrm{Fix}}_R(f)$ (in the last step below), we have

$$\begin{aligned} d_X(f^n\nu, \mu) &\le d_X(f^n\nu, f^n\mu) + d_X(\mu, f^n\mu) \\ &\le d_{f^{-n}(X)}(\nu, \mu) + \sum_{i=0}^{n-1} d_{f^{-i}(X)}(\mu, f\mu) \\ &\le M_1 + nR. \end{aligned}$$

Thus,

$$d_X(\mu_0, f^n\nu) \ge d_X(\mu_0, \mu) - d_X(f^n\nu, \mu) > 2M_1.$$

By Corollary 2.8.4,

$$d_{f^{-n}(X)}(f^n\nu,\mu)\leq 2M_1.$$

As above, we iterate the argument on taking powers of f^n. For all $i\geq 0$, we have

$$\begin{aligned} d_X(f^{(i+1)n}\nu,\mu) &\leq d_X(f^{(i+1)n}\nu,f^n\mu)+d_X(\mu,f^n\mu)\\ &\leq d_{f^{-n}(X)}(f^{in}\nu,\mu)+nR\\ &\leq 2M_1+nR. \end{aligned}$$

This eventually leads to the contradiction

$$\begin{aligned} d_X(\mu_0,\mu) &\leq d_X(\mu_0,\nu)+d_X(\nu,\mu)\\ &\leq M_1+2M_1+nR\\ &\leq 3M_1+nR. \end{aligned}$$

We conclude X and $f^{-n}(X)$ must be disjoint. □

3.4 Second technical lemma

In the following, we prove another technical result, which has a similar conclusion as Lemma 3.3.4, but it is based on different assumptions. Its purpose is for the situation when the main surface $\mathcal{S}$ is a bad domain for an R–fixed point μ of a finite order mapping class f (see Proposition 4.2.1). In this situation, we need to cook up a set of base curves for a new fixed point of f which is closer to the base marking μ_0 in $\mathcal{C}(\mathcal{S})$. Lemma 3.4.1 starts this process by finding a subsurface whose orbit under f are all disjoint. Furthermore, the boundary curves of these subsurfaces form a multicurve which is closer to μ_0 in $\mathcal{C}(\mathcal{S})$ than the base curves of μ.

The proof of Lemma 3.4.1 will be similar to that of Lemma 3.3.4. We will provide the details of the first half of the proof to illustrate the differences and omit the second half.

In the following, let R be any constant and let $\mu\in\widetilde{\mathrm{Fix}}_R(f)$ for a finite order mapping class f. Let $H(\mu_0,f\mu_0)$ and $H(\mu_0,\mu)$ be hierarchies. Let $[v_0,f(v_0)]$ and $[v_0,v]$ be the corresponding main geodesics in $\mathcal{C}(\mathcal{S})$. For any domain $Y\subset\mathcal{S}$, we adopt the notation $[u,v]_Y$ to mean the line segment $\big[\pi_Y(u),\pi_Y(v)\big]$ in $\mathcal{C}(Y)$. For any two sets $A,B\subset\mathcal{C}(\mathcal{S})$, let $\mathrm{Dist}_{\mathcal{S}}(A,B)=\min\limits_{\substack{u\in A\\ v\in B}} d_S(u,v)$.

Lemma 3.4.1. *There exists a constant Δ, depending only on R, such that the following holds. Suppose b is a vertex on $[v_0,v]$ with the property that*

$$\mathrm{Dist}_{\mathcal{S}}\Big(\big[b,f(b)\big],\big[v_0,f(v_0)\big]\Big)\geq 4.$$

Let μ' be a separating marking at b. Then whenever a subsurface $Y\subset\mathcal{S}$ has the property that

$$\mathrm{Dist}_{\mathcal{S}}\Big(\partial Y,\big[b,f(b)\big]\Big)\leq 1 \quad \textit{and} \quad d_Y(\mu',f\mu')>\Delta,$$

then

$$Y,f(Y),\ldots,f^{L_Y}(Y)$$

are all domains for geodesics in $H(\mu_0,\mu)$ and are all mutually disjoint.

Proof. We claim the constant

$$\Delta=(2N+1)M_0+(2N+1)R+10M_1$$

works. Let $Y\subset\mathcal{S}$ satisfy the criteria of the lemma. As in Lemma 3.3.4, we will prove by induction on $f^{-i}(Y)$, for $i=0\ldots,L_Y$.

Let's first show Y supports a geodesic in $H(\mu_0, \mu)$. We will be considering $H(f\mu_0, f\mu)$ in parallel. Note that $f\mu'$ will be a separating marking at $f(b)$ in $H(f\mu_0, f\mu)$. Consider the following pair of quadrilaterals in $\mathcal{C}(Y)$:

$$Q_1 = \big[\mu_0, f\mu_0\big]_Y \cup \big[f\mu_0, f\mu')\big]_Y \cup \big[\mu_0, \mu'\big]_Y \cup \big[\mu', f\mu')\big]_Y,$$

and

$$Q_2 = \big[\mu', f\mu'\big]_Y \cup \big[\mu', \mu\big]_Y \cup \big[\mu, f\mu\big]_Y \cup \big[f\mu', f\mu\big]_Y.$$

By our assumption, $d_Y(\mu', f\mu') > \Delta$. By the triangle inequality, at least one of three other segments of Q_1 is long:

$$\big[\mu_0, \mu'\big]_Y, \quad \big[\mu_0, f\mu_0\big]_Y, \quad \big[f\mu_0, f\mu'\big]_Y.$$

Similarly, at least one of the following segments of Q_2 is long:

$$\big[\mu', \mu\big]_Y, \quad \big[\mu, f\mu)\big]_Y, \quad \big[f\mu', f\mu)\big]_Y.$$

Since $d_Z(\mu, f\mu) \leq R$ for all $Z \subseteq \mathcal{S}$, in Q_2 the situation reduces to either

$$d_Y(\mu', \mu) > \frac{\Delta - R}{2} > NM_0 + NR + 5M_1, \tag{13}$$

or

$$d_Y(f\mu', f\mu) > NM_0 + NR + 5M_1. \tag{14}$$

If (13) holds, then by the fact that μ' is a separating marking at b (Lemma 2.9.2),

$$d_Y(\mu_0, \mu') \leq M_1.$$

Applying the triangle inequality yields

$$\begin{aligned} d_Y(\mu_0, \mu) &\geq d_Y(\mu', \mu) - d_Y(\mu_0, \mu') \\ &> (NM_0 + NR + 5M_1) - M_1. \end{aligned}$$

Therefore (13) implies Y supports a geodesic in H. So we may assume (14) holds.

In Q_1, the assumption on ∂Y forces $\mathrm{Dist}_{\mathcal{S}}\Big(\partial Y, \big[\mu_0, f\mu_0\big]\Big) > 1$. In other words, every vertex in $\big[\mu_0, f\mu_0\big]$ crosses Y. Theorem 2.4.3 applies and $d_Y(\mu_0, f\mu_0) \leq M_0$. The situation is reduced to either

$$d_Y(\mu_0, \mu') > \frac{\Delta - M_0}{2} > NM_0 + NR + 5M_1 \tag{15}$$

or

$$d_Y(f\mu_0, f\mu') > NM_0 + NR + 5M_1. \tag{16}$$

It is not possible for (16) to occur simultaneously as (14), as that would mean both

$$d_Y(f\mu_0, f\mu') > M_1 \quad \text{and} \quad d_Y(f\mu', f\mu) > M_1,$$

violating $f\mu'$ a separating marking. So (15) must hold. As above, we must then have

$$\begin{aligned} d_Y(\mu_0, \mu). &\geq d_Y(\mu_0, \mu') - d_Y(\mu', \mu) \\ &> (NM_0 + NR + 5M_1) - M_1. \end{aligned}$$

Therefore, in all cases, Y must support a geodesic in $H(\mu_0, \mu)$. See Figure 2 for a schematic picture of Q_1 and Q_2. Note that the conclusion of the base case always resulted in

$$d_Y(\mu_0, \mu) > NM_0 + NR + 4M_1. \tag{17}$$



Figure 2: The quadrilaterals Q_1 and Q_2 in $\mathcal{C}(Y)$

By induction,

$$Y, f^{-1}(Y), \ldots, f^{-n+1}(Y)$$

are all domains for geodesics in $H(\mu_0, \mu)$ and are all mutually disjoint. Let's now prove $f^{-n}(Y)$ supports a geodesic in $H(\mu_0, \mu)$. Since

$$\mathrm{Dist}_{\mathcal{S}}\Big(\big[v_0, f(v_0)\big], \big[b, f(b)\big]\Big) \geq 4 \quad \text{and} \quad \mathrm{Dist}_{\mathcal{S}}\Big(\partial Y, \big[b, f(b)\big]\Big) \leq 1,$$

the disjointness condition will imply

$$\mathrm{Dist}_{\mathcal{S}}\Big(\big[v_0, f(v_0)\big], \partial f^{-i}(Y)\Big) \geq 2,$$

for all $i = 0, \ldots, n-1$. By Theorem 2.4.3,

$$d_{f^{-i}(Y)}(\mu_0, f\mu_0) \leq M_0. \tag{18}$$

Coupling this fact with (17) (in the last step below), we have

$$\begin{aligned}
&d_{f^{-n}(Y)}(\mu_0, \mu)\\
&\qquad = d_{f^{-n+1}(Y)}(f\mu_0, f\mu)\\
&\qquad \geq d_{f^{-n+1}(Y)}(\mu_0, \mu) - d_{f^{-n+1}(Y)}(\mu_0, f\mu_0) - d_{f^{-n+1}(Y)}(\mu, f\mu)\\
&\qquad \geq d_{f^{-n+1}(Y)}(\mu_0, \mu) - M_0 - R\\
&\qquad\qquad \vdots\\
&\qquad \geq d_Y(\mu_0, \mu) - nM_0 - nR\\
&\qquad > (N-n)M_0 + (N-n)R + 4M_1.
\end{aligned}$$

Since $N > L_Y \geq n$, the above in particular implies

$$d_{f^{-n}(Y)}(\mu_0, \mu) > M_1 \geq M_2.$$

So $f^{-n}(Y)$ supports a geodesic in $H(\mu_0, \mu)$.

Now let's assume $Y, \ldots, f^{-n+1}(Y)$ are all mutually disjoint. Using the action f, we see that $f^{-n}(Y)$ is disjoint with each $Y, \ldots, f^{-n+1}(Y)$ if and only if $f^{-n}(Y)$ and Y are disjoint. If Y and $f^{-n}(Y)$ are not disjoint, then they are time-ordered in $H(\mu_0, \mu)$. The two different cases of time-ordering of Y and $f^{-n}(Y)$ will both lead to a contradiction. The argument is very similar to the one given in Lemma 3.4.1. We will quickly give the argument in the case that $Y <_t f^{-n}(Y)$ and omit the case the argument in the second case.

Suppose $Y <_t f^{-n}(Y)$. Let ν be a hierarchy marking $H(\mu_0, \mu)$ such that

$$d_Y(\nu, \mu) \leq M_1 \quad \text{and} \quad d_{f^{-n}(Y)}(\mu_0, \nu) \leq M_1, \tag{19}$$

as in Lemma 2.8.4. Then

$$\begin{aligned}
d_Y(\mu_0, f^n\nu) &\leq d_Y(\mu_0, f^n\mu_0) + d_Y(f^n\mu_0, f^n\nu) \\
&\leq \left(\sum_{j=0}^{n-1} d_Y(f^j\mu_0, f^{j+1}\mu_0)\right) + d_{f^{-n}(Y)}(\mu_0, \nu) \\
&\leq \left(\sum_{j=0}^{n-1} d_{f^{-j}(Y)}(\mu_0, f\mu_0)\right) + M_1 \\
&\leq nM_0 + M_1.
\end{aligned}$$

Using (17) and the triangle inequality, we have

$$d_Y(f^n\nu, \mu) \geq d_Y(\mu_0, \mu) - d_Y(\mu_0, f^n\nu) > 2M_1.$$

Therefore, by Corollary 2.8.4,

$$d_{f^{-n}(Y)}(\mu_0, f^n\nu) \leq 2M_1.$$

By considering powers of $f^n\nu$ inductively, we have

$$\begin{aligned}
d_Y(\mu_0, f^{(i+1)n}\nu) &\leq d_Y(\mu_0, f^n\mu_0) + d_Y(f^n\mu_0, f^{(i+1)n}\nu) \\
&\leq nM_0 + d_{f^{-n}(Y)}(\mu_0, f^{in}\nu) \\
&\leq nM_0 + 2M_1
\end{aligned}$$

This is true for every $i \geq 0$. Since $N > L_Y \geq n$, using (19), we have

$$d_Y(\mu_0, \mu) \leq d_Y(\mu_0, \nu) + d_Y(\nu, \mu) \leq nM_0 + 3M_1,$$

contradicting (17). The case of $f^{-n}(Y) <_t Y$ will lead to a similar contradiction. This concludes the proof of the lemma. ☐

4 L.B.C. property for finite order mapping classes

The heart of this section is to prove Theorem C in the introduction, which is restated below. Let R_1 be the fixed constant of Lemma 3.1.4.

Theorem 4.0.2. *There exists a constant $R \geq R_1$, depending only on $\mathcal{S}$, such that any finite order $f \in \mathcal{MCG}(\mathcal{S})$ has a marking $\mu \in \widetilde{\mathrm{Fix}}_R(f)$ such that*

$$d_M(\mu_0, \mu) \prec d_M(\mu_0, f\mu_0).$$

Assuming Theorem 4.0.2, we can derive L.B.C. property for finite order mapping classes by a standard argument, following [BH99]. We first state and prove the following corollary of Theorem 4.0.2, which reduces L.B.C. property for finite order mapping classes to a finite problem.

Corollary 4.0.3. *There exists a finite set $\Gamma \subset \mathcal{MCG}(\mathcal{S})$ such that, for every finite order $f \in \mathcal{MCG}(\mathcal{S})$, there exists $\omega \in \mathcal{MCG}(\mathcal{S})$ such that $\omega^{-1}f\omega \in \Gamma$ and $|\omega| \prec |f|$.*

Proof. By enlarging R if necessary, we may rephrase Theorem 4.0.2 in terms of fixed points: there exists R depending only on $\mathcal{S}$ such that any finite order mapping class f has a marking $\mu \in \mathrm{Fix}_R(f)$ such that

$$d_M(\mu_0, \mu) \prec d_M(\mu_0, f\mu_0). \tag{20}$$

We construct the set Γ as follows. Let D be the diameter of $\mathrm{Mark}(\mathcal{S})$ modulo the action of $\mathcal{MCG}(\mathcal{S})$. ($D$ is finite since the action of $\mathcal{MCG}(\mathcal{S})$ on $\mathrm{Mark}(\mathcal{S})$ is cofinite). Set

$$\Gamma = \{\ g \in \mathcal{MCG}(\mathcal{S}) \ \ : \ \ d_M(\mu_0, g\mu_0) \le 2D + R,\ g \text{ finite order } \}.$$

The action of $\mathcal{MCG}(\mathcal{S})$ on $\mathrm{Mark}(\mathcal{S})$ is proper, thus Γ is a finite set. We show Γ satisfies the other properties as well.

Let $f \in \mathcal{MCG}(\mathcal{S})$ be of finite order. Let μ be a R-fixed point of f closest to μ_0. Since the action of $\mathcal{MCG}(\mathcal{S})$ on $\mathrm{Mark}(\mathcal{S})$ is cofinite, there exists $\omega \in \mathcal{MCG}(\mathcal{S})$ such that $d_M(\omega\mu_0, \mu) \le D$. Then $\omega^{-1} f \omega \in \Gamma$, since

$$\begin{aligned} d_M(\mu_0, \omega^{-1} f \omega \mu_0) &= d_M(\omega\mu_0, f\omega\mu_0) \\ &\le d_M(\omega\mu_0, \mu) + d_M(\mu, f\mu) + d_M(f\mu, f\omega\mu_0) \\ &\le d_M(\omega\mu_0, \mu) + d_M(\mu, f\mu) + d_M(\mu, \omega\mu_0) \\ &\le 2D + R. \end{aligned}$$

Moreover, by (20) we have

$$\begin{aligned} |\omega| &\prec d_M(\mu_0, \omega\mu_0) \\ &\le d_M(\mu_0, \mu) + d_M(\omega\mu_0, \mu) \\ &\prec d_M(\mu_0, f\mu_0) + D \\ &\prec |f|. \end{aligned}$$

□

Corollary 4.0.4 (L.B.C. property for finite order mapping classes)**.** *If $f, g \in \mathcal{MCG}(\mathcal{S})$ are conjugate finite order mapping classes, then there is a conjugating element $\omega \in \mathcal{MCG}(\mathcal{S})$ with*

$$|\omega| \prec |f| + |g|.$$

Proof. Let $\Gamma \subset \mathcal{MCG}(\mathcal{S})$ be the finite set of Theorem 4.0.2. The content of Theorem 4.0.2 is that Γ contains at least one and at most finitely many representatives for each conjugacy class of a finite order mapping class. Furthermore, each finite order f can be conjugated into Γ by a conjugating element whose word length is proportional to $|f|$. The result follows after picking a conjugating element for each pair of elements in Γ of the same conjugacy class. □

The proof of Theorem 4.0.2 will occupy the rest of the section. The main observation is that if $\mu_1 \in \widetilde{\mathrm{Fix}}_{R_1}(f)$ does not have any R_1–bad domains, then μ_1 would satisfy the statement of Theorem 4.0.2. If μ_1 does have a R_1-bad domain Y, then we can construct a marking $\mu_2 \in \widetilde{\mathrm{Fix}}_{R_2}(f)$, where R_2 depends only on R_1, such that $Y \notin \Omega(\mu_2, R_2)$. Ideally, we would like $\Omega(\mu_2, R_2)$ to be strictly smaller than $\Omega(\mu_1, R_1)$, but the situation is a bit more complicated. In trying to improve μ_1 in Y, we may have created new bad domains but we will have control over what they are in relation to Y. We call this the base step of the proof. Although the set of bad domains are not necessarily decreasing, applying the base step in the right way will guarantee a decrease in the complexity of $\Omega(\mu_2, R_2)$ from that of $\Omega(\mu_1, R_1)$. By iterating this process, we produce a sequence of symmetric points for f such that the complexities of the sets of bad domains are monotonically decreasing. This process must stop to produce an R–symmetric point μ for f with no bad domains. Since the maximal complexity of any set of bad domains is the complexity of $\mathcal{S}$, the process of achieving μ terminates after a bounded number of steps. This serves to ensure the constant R will depend only on R_1 and hence only on $\mathcal{S}$.

The rest of the section is organized as follows. We first prove in Proposition 4.1.1 that no bad domain indeed implies Theorem 4.0.2. Then the base step is dealt with in §4.2. There are two propositions, Proposition 4.2.1 and Proposition 4.2.3, associated to the base step, depending on whether the bad domain is the main surface or a proper subsurface. This is where our work in §3 will come in. In §4.3, we explain how to use the base step to reduce complexity of the set of bad domains. The precise statement is Proposition 4.3.1. The section will conclude with Corollary 4.3.2 which makes precise how the process terminates after a bounded number of steps.

4.1 No bad domains

Proposition 4.1.1 (No bad domains)**.** *Let* $\mu \in \widetilde{\mathrm{Fix}}_R(f)$ *where* $R \geq R_1$*. If* R *is a constant depending only on* R_1 *such that* $\Omega(\mu, R) = \emptyset$*, then*

$$d_M(\mu_0, \mu) \prec d_M(\mu_0, f\mu_0).$$

In other words, μ *and* R *satisfy Theorem 4.0.2.*

Proof. The assumption $\Omega(\mu, R) = \emptyset$ means that for every $X \subseteq \mathcal{S}$, there exists i_X such that

$$d_X(\mu_0, \mu) \leq 2N d_{f^{i_X}(X)}(\mu_0, f\mu_0) + NR + \Theta.$$

Let L_0 be the constant of the distance formula, Theorem 2.6.5. Let

$$\Phi = \{ \ X \subseteq \mathcal{S} \ : \ d_X(\mu_0, \mu) \geq 2NL_0 + NR + \Theta \ \},$$

and

$$\Psi = \{ \ Y \subseteq \mathcal{S} \ : \ d_Y(\mu_0, f\mu_0) \geq L_0 \ \}.$$

Then there is a map $\Phi \to \Psi$ sending $X \mapsto f^{i_X}(X)$. This map has multiplicity at most the order of f, which is bounded by N. Therefore,

$$\begin{aligned} d_M(\mu_0, \mu) &\asymp \sum_{X \in \Phi} d_X(\mu_0, \mu) \\ &\prec \sum_{X \in \Phi} d_{f^{i_X}(X)}(\mu_0, f\mu_0) \\ &\leq N \sum_{Y \in \Psi} d_Y(\mu_0, f\mu_0) \\ &\asymp d_M(\mu_0, f\mu_0). \end{aligned}$$

□

4.2 Base step

We are now ready to state and prove the base step of the proof for Theorem 4.0.2. There are two cases to consider, which are Proposition 4.2.1 and Proposition 4.2.3. The proof of Proposition 4.2.1 will be essential for Proposition 4.2.3. Recall the definition of $\xi\big(\Omega(\mu, R)\big)$ as in Definition 3.3.3.

Proposition 4.2.1 (Base Step 1)**.** *Let* $R_I \geq \max\{2\delta + 4, R_1\}$*, and* $\mu_I \in \widetilde{\mathrm{Fix}}_{R_I}(f)$*. If* $\mathcal{S} \in \Omega(\mu_I, R_I)$*, then there exists* R_O*, depending only on* R_I*, and* $\mu_O \in \mathrm{Mark}(\mathcal{S})$*, satisfying the following properties:*

(P1) $\mu_O \in \widetilde{\mathrm{Fix}}_{R_O}(f)$*.*

(P2) $\Omega(\mu_O, R_O) \subsetneqq \Omega(\mu_I, R_I)$*. In addition,* $\mathcal{S} \notin \Omega(\mu_O, R_O)$*, and thus* $\xi(\mu_O, R_O) \lneqq \xi(\mu_I, R_I)$*.*

Proof. Let μ_0 be the base marking in $\mathrm{Mark}(\mathcal{S})$. We have four hierarchies:

$$H(\mu_0, f\mu_0), \quad H(\mu_I, f\mu_I), \quad H(\mu_0, \mu_I), \quad H(f\mu_0, f\mu_I).$$



Figure 3: The quadrilateral Q in $\mathcal{C}(\mathcal{S})$

Consider the four main geodesics corresponding to the four hierarchies, forming a quadrilateral Q in $\mathcal{C}(\mathcal{S})$:

$$\big[v_0, f(v_0)\big], \quad \big[v_I, f(v_I)\big], \quad \big[v_0, v_I\big], \quad \big[f(v_0), f(v_I)\big],$$

where v_0 and v_I are base curves in μ_0 and μ_I, respectively. Our assumption is that $\mathcal{S} \in \Omega(\mu_I, R_I)$, so

$$d_{\mathcal{S}}(v_0, v_I) > d_{\mathcal{S}}\big(v_0, f(v_0)\big) + R_I.$$

Since f acts on $\mathcal{C}(\mathcal{S})$ as an isometry, $d_{\mathcal{S}}(v_0, v_I) = d_{\mathcal{S}}\big(f(v_0), f(v_I)\big)$, so Q is 2δ–thin. The geodesics $[v_I, v_0]$ and $[f(v_I), f(v_0)]$ 2δ–fellow travel for awhile until $[v_I, v_0]$ begins fellow traveling $[v_0, f(v_0)]$. Choose the vertex b on $[v_I, v_0]$ at the junction where this change takes place. After possibly moving b toward v_I, we may assume the following properties for b (see Figure 3):

- $d_{\mathcal{S}}\big(b, f(b)\big) \le 2\delta$, and
- $\mathrm{Dist}_{\mathcal{S}}\Big(\big[b, f(b)\big], \big[v_0, f(v_0)\big]\Big) \ge 4.$

We have

$$d_{\mathcal{S}}(b, v_0) \le d_{\mathcal{S}}\big(v_0, f(v_0)\big) + 2\delta + 4. \tag{21}$$

Now choose the separating marking μ in $H(\mu_0, \mu_I)$ at b. To decide if μ is the right marking for the proposition, consider any $Y \subset \mathcal{S}$ such that

$$\mathrm{Dist}_{\mathcal{S}}\left(\partial Y, [b, f(b)]\right) \le 1. \tag{22}$$

According to Lemma 3.4.1, either $d_Y(\mu, f\mu) \le \Delta$ (see Case I below), where

$$\Delta = (2N+1)M_0 + (2N+1)R_I + 10M_1,$$

or Y and all its orbits under f are disjoint (see Case II).

Case I. If $d_Y(\mu, f\mu) \le \Delta$ for all Y of type (22), then $\mu_O = \mu$ and $R_O = \Delta$ are our desired marking and constant for Proposition 4.2.1. Indeed, the fact that $\mathcal{S} \notin \Omega(\mu_O, R_O)$ follows from (21). Since μ_O comes from a slice in $H(\mu_0, \mu_I)$, we also have

$$d_Z(\mu_0, \mu_O) \le d_Z(\mu_0, \mu_I) + M_3,$$

for all $Z \subset \mathcal{S}$, so (P2) is verified. To see property (P1), we consider $d_Z(\mu_O, f\mu_O)$ for three possibilities of Z.

(a1) If $d_{\mathcal{S}}\big(\partial Z, [b, f(b)]\big) \le 1$, then

$$d_Z(\mu_O, f\mu_O) \le \Delta \le R_O.$$

(a2) If $d_{\mathcal{S}}\big(\partial Z, [b, f(b)]\big) > 1$, then every vertex of $\big[b, f(b)\big]$ cuts Z. By Theorem 2.4.3,

$$d_Z\big(b, f(b)\big) \le M_0 \quad \Longrightarrow \quad d_Y(\mu_O, f\mu_O) \le M_0 + 4.$$

(a3) If $Z = \mathcal{S}$, then by construction, $d_{\mathcal{S}}\big(b, f(b)\big) \le 2\delta$, thus

$$d_{\mathcal{S}}(\mu_O, f\mu_O) \le 2\delta + 4.$$

Case II. On the other hand, if there is some Y of type (22) such that $d_Y(\mu, f\mu) > \Delta$, then Lemma 3.4.1 implies

$$c = \partial Y \cup \partial f(Y) \cup \cdots \cup \partial f^{L_Y}(Y)$$

is a multicurve. In this case, we let μ_O be a marking extension of c relative to μ_I, as in Definition 2.5.6. In particular, $c \subset \text{base}(\mu_O)$. Set

$$R_O = \max\big\{R_I + 2M_3, 4\delta + 9\big\}.$$

We finish the proof by verifying properties (P1) and (P2) for μ_O and R_O. Consider the following analyses.

(b1) If $Z = \mathcal{S}$, then, since both μ_O and $f\mu_O$ contain c as base curves,

$$d_{\mathcal{S}}(\mu_O, f\mu_O) \le d_{\mathcal{S}}(\mu_O, c) + d_{\mathcal{S}}(c, f\mu_O) \le 4.$$

Also, since

$$\mathrm{Dist}_{\mathcal{S}}\Big(\partial Y, \big[b, f(b)\big]\Big) \le 1,$$

we have

$$\begin{aligned} d_{\mathcal{S}}(\mu_0, \mu_O) &\le d_{\mathcal{S}}(v_0, \partial Y) + 4 \\ &\le d_{\mathcal{S}}(v_0, b) + d_{\mathcal{S}}\big(b, f(b)\big) + 5 \\ &\le \Big(d_{\mathcal{S}}\big(v_0, f(v_0)\big) + 2\delta + 4\Big) + 2\delta + 5 \\ &\le d_{\mathcal{S}}\big(v_0, f(v_0)\big) + 4\delta + 9. \end{aligned} \tag{23}$$

(b2) If $Z \ne \mathcal{S}$, but some curve α in c crosses Z, then

$$d_Z(\mu_O, f\mu_O) \le d_Z(\mu_O, \alpha) + d_Z(\alpha, f\mu_O) \le 8.$$

Furthermore, since $f^i(Y)$ all support a geodesic in $H(\mu_0, \mu_I)$, there exists a slice of $H(\mu_0, \mu_I)$ containing α. Therefore,

$$\begin{aligned} d_Z(\mu_0, \mu_O) &\le d_Z(\mu_0, \alpha) + 4 \\ &\le d_Z(\mu_0, \mu_I) + M_3 + 4. \end{aligned}$$

(b3) If $Z \subset \mathcal{S}$ is such that Z is disjoint from c or Z is curve in c, then we are in the situation of Lemma 2.5.7. Note that $f\mu_O$ is a marking extension of c relative to $f\mu_I$.

$$\begin{aligned} d_Z(\mu_O, f\mu_O) &\le d_Z(\mu_O, \mu_I) + d_Z(\mu_I, f\mu_I) + d_Z(f\mu_I, f\mu_O) \\ &\le d_Z(\mu_I, f\mu_I) + 2M_3 \\ &\le R_I + 2M_3, \end{aligned}$$

and

$$d_Z(\mu_0, \mu_O) \le d_Z(\mu_0, \mu_I) + d_Z(\mu_I, \mu_O) \le d_Z(\mu_0, \mu_I) + M_3.$$

Together above we have shown $\mu_O \in \widetilde{\mathrm{Fix}}_{R_O}(f)$. From (23) we have $\mathcal{S} \notin \Omega(\mu_O, R_O)$. Finally, if $Z \in \Omega(\mu_O, R_O)$, then Z must be of case (b2) or (b3). In either case, we have

$$\begin{aligned} d_Z(\mu_0, \mu_I) &\geq d_Z(\mu_0, \mu_O) - (M_3 + 4) \\ &> 2N\Big(\max_i \big\{d_{f^i(Z)}(\mu_0, f\mu_0)\big\}\Big) + NR_O + \Theta - (M_3 + 4) \\ &> 2N\Big(\max_i \big\{d_{f^i(Z)}(\mu_0, f\mu_0)\big\}\Big) + N(R_I + 2M_3) + \Theta - (M_3 + 4) \\ &> 2N\Big(\max_i \big\{d_{f^i(Z)}(\mu_0, f\mu_0)\big\}\Big) + NR_I + \Theta \end{aligned}$$

Therefore, $\Omega(\mu_O, R_O) \subset \Omega(\mu_I, R_I)$, as desired. ☐

Remark 4.2.2. Let μ_O and R_O be the pair satisfying Proposition 4.2.1 coming from either case I or II in the proof. We emphasize that that in establishing the inclusion $\Omega(\mu_O, R_O) \subset \Omega(\mu_I, R_I)$, we essentially proved that for all $Z \subset \mathcal{S}$,

$$d_Z(\mu_0, \mu_O) \leq d_Z(\mu_0, \mu_I) + M_3 + 4.$$

The inclusion was strict since $\mathcal{S} \notin \Omega(\mu_O, R_O)$ by construction.

Proposition 4.2.3 (Base Step 2)**.** *Let R_I and μ_I be as in Proposition 4.2.1. If $\mathcal{S} \notin \Omega(\mu_I, R_I)$ but $\Omega(\mu_I, R_I)$ contains a proper domain $X \subset \mathcal{S}$, then there exists a constant R_O, depending only on R_I, and $\mu_O \in \mathrm{Mark}(\mathcal{S})$, satisfying the following properties:*

- *(Q1) $\mu_O \in \widetilde{\mathrm{Fix}}_{R_O}(f)$.*
- *(Q2) Let $c = \partial X \cup \partial f(X) \cup \cdots \partial f^{L_X}(X)$. Then $c \subseteq \mathrm{base}(\mu_O)$.*
- *(Q3) For all $i = 0, \ldots L_X$, $f^i(X) \notin \Omega(\mu_O, R_O)$.*
- *(Q4) Suppose $Z \in \Omega(\mu_I, R_I)$ has the property that Z interlocks $f^i(X)$. If $X <_t f^{-i}(Z)$ in $H(\mu_0, \mu_I)$, then $Z \notin \Omega(\mu_O, R_O)$.*
- *(Q5) If $Z \in \Omega(\mu_O, R_O)$ but $Z \notin \Omega(\mu_I, R_I)$, then Z must be a subsurface of $f^i(X)$, for some i. In particular, $\xi(Z) < \xi(X)$.*

Remark 4.2.4. We first note that by Lemma 3.3.4,

$$X, f(X), \ldots f^{L_X}(X)$$

are all disjoint, so c is a simplex in $\mathcal{C}(\mathcal{S})$ and property (Q2) makes sense. (Recall that if X is a curve, then $\partial X = X$.)

Secondly, the assumptions in (Q4) also make sense. If $Z \in \Omega(\mu_I, R_I)$ interlocks $f^i(X)$, then X and $f^{-i}(Z)$ interlock by the action of f. Since they both support geodesics in $H(\mu_0, \mu_I)$ (Lemma 3.3.4), they must be time-ordered. The proposition analyzes the case that $X <_t f^{-i}(Z)$.

The point of property (Q3) is that, if X is a bad domain for μ_I, then we can improve μ_I in $X, f(X), \ldots f^{L_X}(X)$ simultaneously. This process also eliminates all bad domains of type specified by (Q4). However, during this process, new bad domains that do not come from bad domains for μ_I may have been created. Property (Q5) puts restrictions on such bad domains, namely, they can only be subsurfaces $f^i(X)$, which have strictly smaller complexity than that of X. If X is a curve, then in particular (Q5) implies such Z cannot exist and thus $\Omega(\mu_O, R_O) \subset \Omega(\mu_I, R_I)$.

Proof. We will construct a marking μ_O containing

$$c = \partial X \cup \cdots \cup \partial f^{L_X}(X).$$

as base curves, guaranteeing (Q2). The situation may seem similar to case II of Proposition 4.2.1, but to ensure (Q3), it will not be enough to construct μ_O by inducing μ_I on each $f^i(X)$. We will in fact need the full work of Proposition 4.2.1 to construct a marking on X. The action of f will then extend this marking to each $f^i(X)$. We will consider two cases, when X is a curve or when X a non-annular subsurface. The two cases are pretty much the same, but for clarity, we treat them separately. After we explain how to construct μ_O and R_O in each case, we will then check that they satisfy the proposition.

First suppose X is a curve. On each component domain of (S, c), we put the induced marking coming from μ_I. To complete this into a marking, we need to pick a transversal to each $f^i(X)$. Much like as in the proof of Proposition 4.2.1, we have a quadrilateral in $\mathcal{C}(X)$ formed by projecting the main geodesics μ_0, $f\mu_0$, μ_I, and $f\mu_I$ to $\mathcal{C}(X)$. Since the pair of geodesics $\big[\mu_0, \mu_I\big]_X$ and $\big[f\mu_0, f\mu_I\big]_X$ 2δ–fellow travel in $\mathcal{C}(X)$, we can find an element $b \in \mathcal{C}(X)$ such that

- $d_X\big(b, f(b)\big) \le 2\delta$.
- $d_X(b, \mu_0) \le d_X(\mu_0, f\mu_0) + 2\delta$.

Now let $f^i(b)$ be the transversal to $f^i(X)$ and let μ_O be the associated clean marking on S. The correct constant will be $R_O = R_I + 2M_3$.

Now suppose X is a non-annular domain. Let $F = f^{L_X+1} : X \to X$ be the first return map of f to X. Set

$$R_I' = NR_I + 2M_3.$$

Let $\nu_0 = \Pi_X(\mu_0)$ and $\nu_I = \Pi_X(\mu_I)$ be respectively the induced markings of μ_0 of μ_I on X. We will regard ν_0 as the base marking in $\mathrm{Mark}(X)$. Since $\mu_I \in \widetilde{\mathrm{Fix}}_{R_I}(f)$, for any $Z \subseteq X$,

$$\begin{aligned} d_Z(\nu_I, F\nu_I) &\le d_Z(\mu_I, F\mu_I) + 2M_3 \\ &\le (L_X + 1)R_I + 2M_3 \\ &< R_I'. \end{aligned}$$

In other words, $\nu_I \in \widetilde{\mathrm{Fix}}_{R_I'}(F)$. Moreover, since $X \in \Omega(\mu_I, R_I)$, we have

$$\begin{aligned} d_X(\nu_0, \nu_I) &\ge d_X(\mu_0, \mu_I) - 2M_3 \\ &> N \max_{0 \le i \le L_X} \big\{ d_{f^i(X)}(\mu_0, f\mu_0) \big\} + NR_I + \Theta - 2M_3 \\ &> \sum_{i=0}^{L_x} d_{f^{-i}(X)}(\mu_0, f\mu_0) + NR_I + \Theta - 2M_3 \\ &= \sum_{i=0}^{L_X} d_X(f^i\mu_0, f^{i+1}\mu_0) + NR_I + \Theta - 2M_3 \\ &\ge d_X(\mu_0, F\mu_0) + NR_I + \Theta - 2M_3 \\ &\ge d_X(\nu_0, F\nu_0) + NR_I + \Theta - 4M_3 \\ &\ge d_X(\nu_0, F\nu_0) + R_I'. \end{aligned}$$

In other words, $X \in \Omega(\nu_I, R_I', F)$. We may apply Proposition 4.2.1, treating X as the whole surface. This gives a marking ν_O on X and a constant $R_O' \ge R_I'$ depending only on R_I' (hence R_I) such that

(P1) For any $Z \subseteq X$,

$$d_Z(\nu_O, F\nu_O) \leq R'_O. \tag{24}$$

(P2) $X \notin \Omega(\nu_O, R'_O, F)$, meaning

$$d_X(\nu_0, \nu_O) \leq d_X(\nu_0, F\nu_0) + R'_O. \tag{25}$$

The action of f induces a marking $f^i\nu_O$ on each $f^i(X)$. We complete

$$c \cup \bigcup_{i=0}^{L_X} f^i \nu_O$$

to a marking μ_O on $\mathcal{S}$ by extending μ_I to the remaining complements and the curves in c. In this case, set $R_O = R'_O + 2M_3$.

Now, for either X is a curve or a non-annular domain, let μ_O and R_O be the appropriate marking and constant. We will show μ_O and R_O satisfy properties (Q1), (Q3), (Q4) and (Q5). Let's consider the following analyses.

(c1) If $Z = \mathcal{S}$, by assumption, $\mathcal{S} \notin \Omega(\mu_I, R_I)$, so

$$d_{\mathcal{S}}(\mu_0, \mu_I) \leq d_{\mathcal{S}}(\mu_0, f\mu_0) + R_I.$$

Since X is a domain of a geodesic in $H(\mu_0, \mu_I)$, we have

$$\begin{aligned} d_{\mathcal{S}}(\mu_0, \mu_O) &\leq d_{\mathcal{S}}(\mu_0, c) + 2 \\ &\leq d_{\mathcal{S}}(\mu_0, \mu_I) + M_3 + 2 \\ &\leq d_{\mathcal{S}}(\mu_0, f\mu_0) + R_I + M_3 + 2 \\ &\leq d_{\mathcal{S}}(\mu_0, f\mu_0) + R_O. \end{aligned}$$

In particular, $S \notin \Omega(\mu_O, R_O)$. As in (b1) of case II in Proposition 4.2.1, we also have

$$d_{\mathcal{S}}(\mu_O, f\mu_O) \leq 4.$$

(c2) If $Z \neq \mathcal{S}$ but some curve of c crosses Z, then the same argument of (b2) of Proposition 4.2.1 applies to give

$$d_Z(\mu_0, \mu_O) \leq d_Z(\mu_0, \mu_I) + M_3 + 4,$$

and

$$d_Z(\mu_O, f\mu_O) \leq 8.$$

(c3) If Z is a subsurface of some component domain of $(\mathcal{S}, c)$ on which μ_O is induced from μ_I (this includes the possibility that Z is a curve in c when X is not a curve), then, as in (b3) of Proposition 4.2.1,

$$d_Z(\mu_0, \mu_O) \leq d_Z(\mu_0, \mu_I) + M_3,$$

and

$$d_Z(\mu_O, f\mu_O) \leq R_I + 2M_3 \leq R_O.$$

(c4) If X is a curve, then by construction

$$\begin{aligned} d_X(\mu_0, \mu_O) &\leq d_X(\mu_0, b) + M_3 \\ &\leq d_X(\mu_0, f\mu_0) + 2\delta + M_3 \\ &\leq d_X(\mu_0, f\mu_0) + R_O, \end{aligned}$$

and

$$\begin{aligned}
d_X(\mu_O, f\mu_O) &\leq d_X(b, f(b)) + 2M_3 \\
&\leq 2\delta + 2M_3 \\
&\leq R_O.
\end{aligned}$$

If X is non-annular, and $Z \subseteq X$, then it follows from (24) that

$$\begin{aligned}
d_Z(\mu_O, f\mu_O) &\leq d_Z(\nu_O, F\nu_O) + 2M_3 \\
&\leq R'_O + 2M_3 \\
&= R_O.
\end{aligned}$$

Finally, (25) yields

$$\begin{aligned}
d_X(\mu_0, \mu_O) &\leq d_X(\nu_0, \nu_O) + 2M_3 \\
&\leq d_X(\nu_0, F\nu_0) + R'_O + 2M_3 \\
&\leq d_X(\mu_0, F\mu_0) + R'_O + 4M_3 \\
&\leq N \max_{0 \leq i \leq L_X} \big\{ d_{f^i(X)}(\mu_0, f\mu_0) \big\} + NR_O + \Theta. \qquad (26)
\end{aligned}$$

One consequence here is that, whether or not X is a curve, $X \notin \Omega(\mu_O, R_O)$.

(c5) If X is a curve and $i \neq 0$, then both μ_O and $f\mu_O$ are close to $f^i(b)$ in $\mathcal{C}\big(f^i(X)\big)$, thus

$$d_{f^i(X)}(\mu_O, f\mu_O) \leq d_{f^i(X)}(f^i(b), f^i(b)) + 2M_3 \leq R_O.$$

Furthermore,

$$\begin{aligned}
d_{f^i(X)}(\mu_0, \mu_O) &\leq d_{f^i(X)}\big(\mu_0, f^i(b)\big) + M_3 \\
&\leq d_X(f^{-i}\mu_0, b) + M_3 \\
&\leq d_X(\mu_0, b) + d_X(\mu_0, f^{-i}(\mu_0)) + M_3 \\
&\leq d_X(\mu_0, b) + \sum_{j=0}^{i-1} d_X(f^{-j}\mu_0, f^{-(j+1)}\mu_0) + M_3 \\
&= d_X(\mu_0, f\mu_0) + 2\delta + \sum_{j=0}^{i-1} d_{f^{j+1}(X)}(\mu_0, f\mu_0) + M_3 \\
&= \sum_{j=0}^{i} d_{f^{j+1}(X)}(\mu_0, f\mu_0) + 2\delta + M_3 \\
&\leq N \max_{0 \leq j \leq L_X} \big\{ d_{f^j(X)}(\mu_0, f\mu_0) \big\} + R_O.
\end{aligned}$$

If X is non-annular, and $Z \subseteq f^i(X)$, $i \neq 0$, then

$$d_Z(\mu_O, f\mu_O) = d_Z(f^i\nu_O, f^i\nu_O) + 2M_3 \leq R_O.$$

For $f^i(X)$, $i \neq 0$, then

$$\begin{aligned}
d_{f^i(X)}(\mu_0, \mu_O) &\leq d_{f^i(X)}(\mu_0, f^i\nu_O) + M_3 \\
&= d_X(f^{-i}\mu_0, \nu_O) + M_3 \\
&\leq d_X(\mu_0, \nu_O) + \sum_{j=0}^{i-1} d_X(f^{-j}\mu_0, f^{-(j+1)}\mu_0) \\
\text{By (26)} \quad &\leq 2N \max_{0\leq j\leq L_X} \left\{ d_{f^j(X)}(\mu_0, f\mu_0) \right\} + NR_O + \Theta
\end{aligned}$$

An consequence here is that $f^i(X) \notin \Omega(\mu_O, R_O)$.

Together from (c1) to (c5), we have shown that $\mu_O \in \widetilde{\mathrm{Fix}}_{R_O}(f)$. Property (Q3) is verified in cases (c4) and (c5). To see (Q5), if $Z \in \Omega(\mu_O, R_O)$, then Z is either of case (c2), (c3), or $Z \subsetneqq f^i(X)$, for some $i = 0, \ldots, L_X$. In (c2) or (c3), since

$$d_Z(\mu_0, \mu_O) \leq d_Z(\mu_0, \mu_I) + M_3 + 4,$$

it follows that (see Remark 4.2.2)

$$Z \in \Omega(\mu_O, R_O) \quad \Longrightarrow \quad Z \in \Omega(\mu_I, R_I).$$

Too see (Q4), we use Lemma 2.8.2 on the assumption $X <_t f^{-i}(Z)$ to obtain

$$d_{f^{-i}(Z)}(\mu_0, \partial X) \leq M_1.$$

Since μ_O contains $c = \bigcup_j \partial f^j(X)$ as base curves, we have

$$\begin{aligned}
d_Z(\mu_0, \mu_O) &= d_{f^{-i}(Z)}(f^{-i}\mu_0, f^{-i}\mu_O) \\
&\leq d_{f^{-i}(Z)}(f^{-i}\mu_0, \mu_0) + d_{f^{-i}(Z)}(\mu_0, f^{-i}\mu_O) \\
&\leq N \max_{0\leq i\leq L_Z} \left\{ d_{f^i(Z)}(\mu_0, f\mu_0) \right\} + d_{f^{-i}(Z)}(\mu_0, \partial X) + M_3 \\
&\leq N \max_{0\leq i\leq L_Z} \left\{ d_{f^i(Z)}(\mu_0, f\mu_0) \right\} + M_1 + M_3
\end{aligned}$$

Therefore, $Z \notin \Omega(\mu_O, R_O)$. This concludes (Q4) and the proof. □

4.3 Reducing complexity

In this section, we show how to use Proposition 4.2.1 and Proposition 4.2.3 to construct R and μ for Theorem 4.0.2.

Proposition 4.3.1 (Reducing complexity)**.** *Let $f \in \mathcal{MCG}(\mathcal{S})$ be of finite order. Let R_I and μ_I be as in Proposition 4.2.1. Suppose $\Omega(\mu_I, R_I) \neq \emptyset$. There exists R_O, depending only on R_I, and $\mu_O \in \widetilde{\mathrm{Fix}}_{R_O}(f)$ such that*

$$\xi(\mu_O, R_O) \lneqq \xi(\mu_I, R_I).$$

Proof. From now on, a pair (μ, R) will always mean $\mu \in \widetilde{\mathrm{Fix}}_R(f)$.

If $\mathcal{S} \in \Omega(\mu_I, R_I)$, then Proposition 4.2.1 produces (μ_O, R_O) such that R_O depends only on R_I and $\mathcal{S} \notin \Omega(\mu_O, R_O)$, hence $\xi(\mu_O, R_O) < \xi(\mathcal{S}) = \xi(\mu_I, R_I)$.

Now suppose $\mathcal{S} \notin \Omega(\mu_I, R_I)$. Choose a maximal element $X_1 \in \Omega(\mu_I, R_I)$. This in particular means X_1 has maximum complexity over all elements of $\Omega(\mu_I, R_I)$. Set

$$\mathcal{U}_1 = X_1 \cup \cdots \cup f^{L_{X_1}}(X_1).$$

We will say a domain $Z \subset \mathcal{S}$ is *supported* on $\mathcal{S} \setminus \mathcal{U}_1$ if Z lies in some component $\mathcal{S} \setminus \mathcal{U}_1$. Note that in the case that X_1 is not a curve, Z can be a boundary curve of some $f^j(X_1)$. Consider the maximal complexity of the elements in $\Omega(\mu_I, R_I)$ supported on $\mathcal{S} \setminus \mathcal{U}_1$. If this complexity is strictly less than $\xi(X_1)$, then we stop. If this complexity is not strictly less than $\xi(X_1)$, then we may choose X_2 of maximal order in $\Omega(\mu_I, R_I)$ supported on $\mathcal{S} \setminus \mathcal{U}_1$ such that $\xi(X_2) = \xi(X_1)$. Set $\mathcal{U}_2$ and c_2 as above. In this case, it makes sense to say $\mathcal{U}_1$ and $\mathcal{U}_1$ are disjoint. Now we repeat this process by considering the maximum complexity of the elements in $\Omega(\mu_I, R_I)$ supported on $\mathcal{S} \setminus (\mathcal{U}_1 \cup \mathcal{U}_2)$. Continuing this way, we eventually *exhaust* $\mathcal{S}$ by a sequence

$$\mathcal{U}_1, \mathcal{U}_2, \ldots, \mathcal{U}_n$$

in the following sense:

- Each $\mathcal{U}_i$ is a disjoint union of subsurfaces of type
$$\mathcal{U}_i = X_i \cup \cdots \cup f^{L_{X_i}}(X_i),$$
where $\xi(X_i) = \xi(X_1)$, for all i.
- $\mathcal{U}_i$ and $\mathcal{U}_j$ are disjoint, for all $i \neq j$.
- The maximal complexity of the bad domains in $\Omega(\mu_I, R_I)$ supported on
$$\mathcal{S} \setminus (\mathcal{U}_1 \cup \cdots \cup \mathcal{U}_n)$$
is strictly less than $\xi(X_1)$.

Note that the exhaustion sequence has length n which is bounded uniformly by a constant depending only on $\mathcal{S}$. Denote by

$$c_i = \partial X_i \cup \cdots \cup \partial f^{L_{X_i}}(X_i).$$

Since the $\mathcal{U}_i$'s are disjoint, one can apply the proof of Proposition 4.2.3 to them simultaneously to construct a pair (μ_O, R_O) such that, μ_O contains $c_1 \cup \cdots \cup c_n$ as base curves, R_O depends only on R_I, and appropriate analogies to properties (Q3), (Q4), and (Q5) hold for all $\mathcal{U}_i$. In particular, the collection of bad domains in $\Omega(\mu_O, R_O)$ are of the following type. If $Z \in \Omega(\mu_O, R_O)$, then either

(i) Z intersects some curve in c_i for some i. This is case (c2) in the proof of Proposition 4.2.3.

(ii) Z is supported on
$$\mathcal{S} \setminus (\mathcal{U}_1 \cup \cdots \cup \mathcal{U}_n).$$
This is case (c3) in the proof of Proposition 4.2.3

(iii) $Z \subsetneq f^j(X_i)$, for some i and j.

Immediately, case (iii) has $\xi(Z) < \xi(X_1)$. Recall that for either case (i) or (ii),

$$Z \in \Omega(\mu_O, R_O) \qquad \Longrightarrow \qquad Z \in \Omega(\mu_I, R_I).$$

Since $\mathcal{S}$ is exhausted by assumption, case (ii) also means $\xi(Z) < \xi(X_1)$. Lastly, suppose Z is of case (i). Choose the minimal index i such that Z intersects a curve in c_i. In other words, Z is supported on

$$\mathcal{S} \setminus (\mathcal{U}_1 \cup \cdots \cup \mathcal{U}_{i-1}),$$

and Z interlocks $f^j(X_i)$, for some j. Our choice of X_i has maximal order among the bad domains supported on

$$\mathcal{S} \setminus (\mathcal{U}_1 \cup \cdots \cup \mathcal{U}_{i-1}).$$

Therefore, if $\xi(Z) = \xi(X_1) = \xi(X_i)$, then $X_i <_t f^{-j}(Z)$ in $H(\mu_0, \mu_I)$. Property (Q5) of Proposition 4.2.3 guarantees that such domains do not appear in $\Omega(\mu_O, R_O)$. Thus, any Z of case (i) must also have $\xi(Z) < \xi(X)$. □

Let R_1 be the minimal constant satisfying Lemma 3.1.4.

Corollary 4.3.2 (Termination)**.** *There exists $R \geq R_1$ depending only on $\mathcal{S}$ such that any finite order $f \in \mathcal{MCG}(\mathcal{S})$ has a $\mu \in \widetilde{\mathrm{Fix}}_R(f)$ satisfying $\Omega(\mu, R) = \emptyset$.*

Proof. Let $\mu_1 \in \widetilde{\mathrm{Fix}}_{R_1}(f)$. If $\Omega(\mu_1, R_1) = \emptyset$, then we are done. If not, then applying Proposition 4.3.1 iteratively yields a sequence of pairs (μ_1, R_1), (μ_2, R_2), ... such that

- $\mu_{i+1} \in \widetilde{\mathrm{Fix}}_{R_{i+1}}(f)$, where R_{i+1} depends only on R_i.
- $\xi_{i+1} \lneqq \xi_i$, where $\xi_i = \xi(\mu_i, R_i)$.

Since ξ_i corresponds to the maximum complexity over elements in $\Omega(\mu_i, R_i)$, and that ξ_i's are strictly decreasing, we must eventually reach a pair (μ_n, R_n) for which $\xi_n = -2$, i.e. $\Omega(\mu_n, R_n) = \emptyset$. Moreover, since elements of $\Omega(\mu_i, R_i)$ come from subsurfaces of $\mathcal{S}$, $\xi_1 \leq \xi(\mathcal{S}) = 3g - 3 + b$. This gives a bound on $n \leq 3g - 1 + b$, and therefore R_n depends only on $\mathcal{S}$. ☐

This concludes the proof of Theorem 4.0.2.

5 L.B.C. property for reducible mapping classes

In this section, we prove L.B.C. property for (infinite-order) reducible elements of $\mathcal{MCG}(\mathcal{S})$. We would like to use an induction argument on subsurfaces. To do so, we make the following observation.

Let $f \in \mathcal{MCG}(\mathcal{S})$ be reducible with canonical reducing system σ (see §2.2.3). If $\mu \in \mathrm{Mark}(\mathcal{S})$ is a marking containing σ as base curves, then so does $f\mu$. This means that any hierarchy $H(\mu, f\mu)$ decomposes into geodesics supported on component domains of $(\mathcal{S}, \sigma)$. For such a marking μ, we can control each component domain independently, allowing the arguments of Theorem 2.2.4 and Corollary 4.0.4 to pass through to subsurfaces. This inspires the definition of a good marking for f (Definition 5.0.3). We construct a finite collection of good markings and prove Theorem E of the introduction. The induction argument on subsurfaces using good markings appears in Proposition 5.0.5. Finally, Corollary 5.0.6 combines the finiteness and the induction argument to finish the proof of L.B.C. property for reducible elements of $\mathcal{MCG}(\mathcal{S})$.

Definition 5.0.3 (Good marking)**.** Let $f \in \mathcal{MCG}(\mathcal{S})$ be an infinite-order reducible element and let σ be its associated canonical reducing system. We say a marking $\mu \in \mathrm{Mark}(\mathcal{S})$ is a *good marking* for f if $\sigma \subseteq \mathrm{base}(\mu)$.

Up to homeomorphisms of $\mathcal{S}$, there are only finitely many multicurves on $\mathcal{S}$. If $f \in \mathcal{MCG}(\mathcal{S})$ is reducible and $f = \omega^{-1}gh$, then $\omega(\sigma_f) = \sigma_g$, where σ_f and σ_g are canonical reducing system for f and g respectively. We fix a representative for each homeomorphism type of a multicurve. Further, for each representative multicurve σ, we complete σ into a finite set of markings representing each homeomorphism type of a marking containing σ as base curves. Let $\mathcal{M}$ the collection of all such *representative markings*. The following is Theorem E of the introduction.

Theorem 5.0.4. *There exists $a \in \mathcal{MCG}(\mathcal{S})$ such that $a^{-1}fa$ has a good marking in $\mathcal{M}$, and*

$$d_M(\mu_0, a\mu_0) \prec d_M(\mu_0, f\mu_0).$$

Furthermore, if f and g are conjugate, then we may choose the same good marking in $\mathcal{M}$ for both of them.

Before proving Theorem 5.0.4, we set up some notations. Fix a multicurve σ on $\mathcal{S}$. Let $\mathrm{Stab}(\sigma) < \mathcal{MCG}(\mathcal{S})$ be the subgroup stabilizing σ as a set:

$$\mathrm{Stab}(\sigma) : \{\ h \in \mathcal{MCG}(\mathcal{S}) \ \ : h(\sigma) = \sigma\ \}.$$

The action of $\mathrm{Stab}(\sigma)$ on the complementary components of $\mathcal{S} \setminus \sigma$ induces an exact sequence:

$$1 \longrightarrow \mathrm{Stab}_0(\sigma) \longrightarrow \mathrm{Stab}(\sigma) \xrightarrow{\pi} \text{Finite Group} \longrightarrow 1. \tag{27}$$

Consider the kernel $\mathrm{Stab}_0(\sigma)$ of the above sequence. If $f \in \mathrm{Stab}_0(\sigma)$, then f acts on each complementary component $Y \in \mathcal{S} \setminus \sigma$. In other words, f defines an element $f|_Y \in \mathcal{MCG}_0(Y)$ for each $Y \in \mathcal{S} \setminus \sigma$, where $\mathcal{MCG}_0(Y) \subset \mathcal{MCG}(Y)$ is the subgroup fixing ∂Y. We have an exact sequence:

$$1 \longrightarrow T_\sigma \longrightarrow \mathrm{Stab}_0(\sigma) \longrightarrow \prod_{Y \in \mathcal{S} \setminus \sigma} \mathcal{MCG}_0(Y) \longrightarrow 1, \tag{28}$$

where T_σ is a free abelian group with basis the Dehn twists along curves in σ. By the classification theorem, $f|_Y$ is either pseudo-Anosov or has finite order. We say an element $f \in \mathrm{Stab}_0(\sigma)$ is *pure* if $f|_Y$ is either pseudo-Anosov or the identity on Y. The order of the finite group in (27) is bounded by a constant N depending only on $\mathcal{S}$. Thus, for any σ and any $f \in \mathrm{Stab}(\sigma)$, $f^N \in \mathrm{Stab}_0(\sigma)$. Moreover, since there are only finitely many subsurfaces of $\mathcal{S}$ up to homeomorphism, one can choose a constant for Lemma 3.1.1 which works for $\mathcal{S}$ and all subsurfaces of $\mathcal{S}$. Thus, there exists some universal power $N = N(S)$ depending only $\mathcal{S}$, such that for any reducible mapping element $f \in \mathcal{MCG}(\mathcal{S})$, f^N is pure.

We can characterize the canonical reducing system for a reducible mapping class as follows. Suppose $f \in \mathrm{Stab}(\sigma)$. Let $g = f^N$ be pure. Then $\sigma = \sigma_f$ is the canonical reducing system for f if for any $\alpha \in \sigma$, at least one of the following holds.

(H1) There exists $Y \in \mathcal{S} \setminus \sigma$ such that α is a boundary component of Y and $g|_Y$ is pseudo-Anosov on Y.

(H2) There exists a subsurface $Z \subset \mathcal{S}$ such that $g|_Z$ is a non-zero power of a Dehn twist along α.

To see this, let $\alpha \in \sigma$ be such that condition (H1) does not hold. Then α must bound two (not necessarily distinct) components X and Y of $\mathcal{S} \setminus \sigma$ such that $g|_X$ and $g|_Y$ are both identity. In this case, let $Z = X \cup Y \cup \alpha$. Then $g|_Z$ is a non-zero power of Dehn twist along α. Otherwise, the first return map of f to Z is of finite order, and one can thus obtain a smaller reducing system for f by removing α, contradicting minimality of σ. Note that this proof also implies that if σ is a canonical reducing system for f, then σ is also the canonical reducing system for any power of f.

Property (H2) implies, for any $n \in \mathbb{N}$ and any $v \in \mathcal{C}(\alpha)$,

$$d_\alpha\big(v, g^n(v)\big) \geq |n|.$$

Compare this with Lemma 2.3.2. Since there are only finitely many domains of $\mathcal{S}$ up to homeomorphism, we can choose N_0 depending only on $\mathcal{S}$ such that the following hold. Let M_2 be the constant of Lemma 2.6.3. For any multicurve σ and any $g \in \mathrm{Stab}_0(\sigma)$, let Y be either a component of $\mathcal{S} \setminus \sigma$ on which g is pseudo-Anosov, or Y is a curve in σ such that property (H2) holds, then for any $n \geq N_0$ and any $v \in \mathcal{C}(Y)$,

$$d_Y\big(v, g^n(v)\big) \geq M_2.$$

For any $g \in \mathrm{Stab}_0(\sigma)$, Equation (2) has the following consequence. If $\mu \in \mathrm{Mark}(\mathcal{S})$ is a good marking for g, then

$$d_M(\mu, g\mu) \asymp \sum_{X \in \mathcal{S} \setminus \sigma} d_{\mathrm{Mark}(X)}(\mu, g|_X \mu) + \sum_{\alpha \in \sigma} d_\alpha(\mu, g\mu), \tag{29}$$

where

$$d_{\mathrm{Mark}(X)}(\mu, g|_X\mu) := d_{\mathrm{Mark}(X)}\big(\Pi_X(\mu), g|_X\Pi_X(\mu)\big) \asymp d_{\mathrm{Mark}(X)}\big(\Pi_X(\mu), \Pi_X(g\mu)\big).$$

We will say an element $g \in \mathrm{Stab}_0(\sigma)$ *does not twist* along $\alpha \in \sigma$ if for any $v \in \mathcal{C}(\alpha)$,

$$\lim_{n\to\infty} \frac{d_\alpha(v, g^n(v))}{n} = 0.$$

In this case, $d_\alpha(\mu, g\mu) \prec 1$, and one can ignore the second summand on the right hand side of Equation (29).

Proof of Theorem 5.0.4. The set $\mathcal{M}$ is finite and each conjugacy class of $\mathcal{MCG}(\mathcal{S})$ has a good marking in $\mathcal{M}$ by construction. Let

$$C = \max\{\ d_M(\mu_0, \mu)\ \ :\ \ \mu \in \mathcal{M}\ \}.$$

Let $f \in \mathcal{MCG}(\mathcal{S})$ be reducible with with canonical reducing system σ_f. Let σ be the representative multicurve for σ_f and let $\mathcal{M}(\sigma) \subset \mathcal{M}$ be the subset of markings containing σ as base curves. Set $F = f^{N_0}$. Let $\mu \in \mathcal{M}$. Any curve $\alpha \in \sigma_f$ is either a domain for a geodesic in $H(\mu, F\mu)$ or is a boundary curve of a domain for a geodesic in $H(\mu, F\mu)$. Let μ' be a marking extension of σ_f relative to $F\mu$. Our construction of μ' has the property that, for any domain $Y \subseteq \mathcal{S}$, the projection of μ' to Y is uniformly close to the geodesic connecting μ and $F\mu$ in $\mathcal{C}(Y)$. Since this is true for any domain Y, we obtain the following:

$$d_M(\mu, \mu') \prec d_M(\mu, F\mu) \prec d_M(\mu, f\mu).$$

Now choose a marking $\mu'' \in \mathcal{M}(\sigma)$ such that there exist $a \in \mathcal{MCG}(\mathcal{S})$ with $a(\mu'') = \mu'$. By construction, $a^{-1}fa$ has canonical reducing system $a^{-1}(\sigma_f) = \sigma \in\subseteq \mathrm{base}(\mu'')$. We also have

$$\begin{aligned} d_M(\mu_0, a\mu_0) &\le d_M(\mu_0, a\mu'') + d_M(a\mu'', a\mu_0) \le d_M(\mu_0, \mu') + C \\ &\le d_M(\mu_0, \mu) + d_M(\mu, \mu') \prec d_M(\mu, f\mu) \\ &\le d_M(\mu, \mu_0) + d_M(\mu_0, f\mu_0) + d_M(f\mu_0, f\mu) \\ &\prec d_M(\mu_0, f\mu_0). \end{aligned}$$

If $g \in \mathcal{MCG}(\mathcal{S})$ is conjugate to f, then σ would also be the representative multicurve for σ_g. Our construction produces an element $b \in \mathcal{MCG}(\mathcal{S})$ such that $b^{-1}gb$ has a good marking in $\mathcal{M}(\sigma)$ and $d_M(\mu_0, b\mu_0) \prec d_M(\mu_0, g\mu_0)$. Since any marking in $\mathcal{M}(\sigma)$ is a good marking for $b^{-1}gb$, including μ'', and $\mathcal{M}(\sigma)$ is a finite set, the second statement follows. ☐

Proposition 5.0.5. *Suppose $f, g \in \mathcal{MCG}(\mathcal{S})$ are two conjugate infinite-order reducible mapping classes with the same canonical reducing system σ. Let μ be a good marking for f and g. Then there exist a constant K_μ and $\omega \in \mathcal{MCG}(\mathcal{S})$ such that ω is a conjugator for f and g, and*

$$d_M(\mu, \omega\mu) \prec K_\mu\big(d_M(\mu, f\mu) + d_M(\mu, g\mu)\big).$$

Proof. Elements of $\mathrm{Stab}_0(\sigma)$ are easier to handle, but a conjugator for f^n and g^n is not a conjugator for f and g. Thus we cannot apply the results of Corollary 4.0.4 and Theorem 2.2.4, to powers of f and g. We must deal with the issue of permuting subsurfaces in the proof of Proposition 5.0.5. Fix a finite collection $\mathcal{P} \subset \mathrm{Stab}(\sigma)$ such that $\pi(\mathcal{P})$ in the exact sequence (27) is onto. Let

$$P = \max\{\ d_M(\mu, a\mu)\ \ :\ \ a \in \mathcal{P}\ \}.$$

Choose $a \in \mathcal{P}$ such that f and $g' = aga^{-1}$ have the following properties:

(i) $\pi(f) = \pi(g')$.

(ii) For any $X \in \mathcal{S} \setminus \sigma$, the first return map to X of f and g' are conjugate.

If the proposition holds for f and g', say, there exist K_μ and $\omega \in \mathcal{MCG}(\mathcal{S})$ such that $f\omega = \omega g'$ and

$$d_M(\mu, \omega'\mu) \prec K_\mu\big(d_M(\mu, f\mu) + d_M(\mu, g'\mu)\big),$$

then ωa and $K_\mu + 2PK_\mu + P$ verify the proposition for f and g. Clearly, ωa is a conjugator for f and g. It remains to check:

$$\begin{aligned} d_M(\mu, \omega a\mu) &\leq d_M(\mu, \omega\mu) + d_M(\omega\mu, \omega a\mu) \\ &\leq d_M(\mu, \omega\mu) + d_M(\mu, a\mu) \\ &\prec K_\mu\big(d_M(\mu, f\mu) + d_M(\mu, aga^{-1}\mu)\big) + P \\ &\leq K_\mu\big(d_M(\mu, f\mu) + d_M(\mu, g\mu) + 2P\big) + P. \end{aligned}$$

Thus, we may assume f and g already satisfy properties (i) and (ii) above. We will find a conjugator $\omega \in \mathrm{Stab}_0(\sigma)$ for f and g. To do this, we will use properties (i) and (ii) and the induction hypothesis to build a conjugating element $\omega_Y \in \mathcal{MCG}_0(Y)$ for each complementary component $Y \in \mathcal{S} \setminus \sigma$. Then we will choose an appropriate lift $\omega \in \mathrm{Stab}_0(Y)$.

Decompose the complementary components of $\mathcal{S} \setminus \sigma$ into orbits under the action f. Pick a representative from each orbit. Let X_1 be one such representative and consider the sequence of distinct complementary subsurfaces of $\mathcal{S} \setminus \sigma$

$$X_1, X_2, \ldots, X_n$$

such that $f(X_i) = X_{i+1}$ and $g(X_i) = X_{i+1}$, for $i = 1, \ldots, n$ and $X_{n+1} = X_1$. Note that $n < N_0$. Set $f_i = f|_{X_i} : X_i \to X_{i+1}$, and similarly for g_i. Set the first return maps $F = f^{n+1}|_{X_1} \in \mathcal{MCG}(X_1)$ and $G = g^{n+1}|_{X_1} \in \mathcal{MCG}(X_1)$. The assumption is that F and G are conjugate in $\mathcal{MCG}(X_1)$. By Theorem 2.2.2, F and G are either pseudo-Anosov or have finite order on X_1. Letting $\nu_1 = \Pi_{X_1}(\mu)$ be the induced marking on X_1, it follows from results of Corollary 4.0.4 and Theorem 2.2.4 that there exist $\omega_1 \in \mathcal{MCG}_0(X_1)$ and $K_1 = K_1(\nu_1, X_1)$ such that $F\omega_1 = \omega_1 G$, and

$$\begin{aligned} d_{\mathrm{Mark}(X_1)}(\nu_1, \omega_1\nu_1) &\leq K_1\big(d_{\mathrm{Mark}(X_1)}(\nu_1, F\nu_1) + d_{\mathrm{Mark}(X_1)}(\nu_1, G\nu_1)\big) \\ &\prec K_1\big(d_M(\mu, F\mu) + d_M(\mu, G\mu)\big) \qquad \text{(By (2))} \\ &\prec K_1\big(d_M(\mu, f^n\mu) + d_M(\mu, g^n\mu)\big) \\ &\prec K_1\big(d_M(\mu, f\mu) + d_M(\mu, g\mu)\big) \end{aligned}$$

Using f and g, we construct for each X_i an element $\omega_i \in \mathcal{MCG}_0(X_i)$ such that $f_i\omega_i = \omega_{i+1}g_i$, for $i = 1, \ldots n$ and $n+1 = 1$. The element $\omega_1 \in \mathcal{MCG}_0(X_1)$ is defined. For each $i = 1, \ldots n$, set

$$\omega_{i+1} = f_i \cdots f_1\omega_1 g_1^{-1} \cdots g_i^{-1}.$$

In particular, $\omega_{n+1} = F\omega_1 G^{-1} = \omega_1$. Let $\nu_i = \Pi_{X_i}(\mu)$. We have

$$\begin{aligned} d_{\mathrm{Mark}(X_i)}(\nu_i, \omega_{i+1}\nu_i) &= d_{\mathrm{Mark}(X_i)}(\nu_i, f_i \cdots f_1\omega_1 g_1^{-1} \cdots g_i^{-1}\nu_i) \\ &= d_{\mathrm{Mark}(X_i)}(\nu_i, f^i|_{X_1}\omega_1 g^{-i}|_{X_i}\nu_i) \\ &= d_{\mathrm{Mark}(X_1)}(f^{-i}|_{X_i}\nu_i, \omega_1 g^{-i}|_{X_i}\nu_i) \\ &\leq d_{\mathrm{Mark}(X_1)}(f^{-i}|_{X_i}\nu_i, \nu_1) + d_{\mathrm{Mark}(X_1)}(\nu_1, \omega_1 g^{-i}|_{X_i}\nu_i) \\ &= d_{\mathrm{Mark}(X_1)}(f^{-i}|_{X_i}\nu_i, \nu_1) + d_{\mathrm{Mark}(X_1)}(\omega_1^{-1}\nu_1, g^{-i}|_{X_i}\nu_i) \\ &\leq d_{\mathrm{Mark}(X_1)}(f^{-i}|_{X_i}\nu_i, \nu_1) + d_{\mathrm{Mark}(X_1)}(\omega_1^{-1}\nu_1, \nu_1) + d_{\mathrm{Mark}(X_1)}(\nu_1, g^{-i}\nu_i) \\ &\prec d_M(\mu, f^i\mu) + d_M(\mu, g^i\mu) + d_{\mathrm{Mark}(X_1)}(\nu_1, \omega_1\nu_1) \\ &\prec K_1\big(d_M(\mu, f\mu) + d_M(\mu, g\mu)\big) \end{aligned}$$

We do this for each orbit of complementary subsurfaces in $\mathcal{S} \setminus \sigma$, building for each $Y \in \mathcal{S} \setminus \sigma$ an element $\omega_Y \in \mathcal{MCG}_0(Y)$. Consider any element $\omega \in \mathrm{Stab}_0(\sigma)$ such that $\omega|_Y = \omega_Y$. Since twisting commute, any ω will satisfy $f\omega = \omega g$ by construction. Thus, we can choose a lift ω which does not twist along any curves in σ. Let $\{K_i\}$ be the constants associated to each orbit and let $K_\mu = \max\{K_i\}$. Using previous work and Equation (29), we obtain

$$\begin{aligned} d_M(\mu, \omega\mu) &\asymp \sum_{X \in \mathcal{S}\setminus\sigma} d_{\mathrm{Mark}(X)}(\mu, \omega_X \mu) + \sum_{\alpha \in \sigma} d_\alpha(\mu, \omega\mu) \\ &\prec \sum_{X \in \mathcal{S}\setminus\sigma} K_\mu\big(d_M(\mu, f\mu) + d_M(\mu, g\mu)\big) \\ &\prec K_\mu\big(d_M(\mu, f\mu) + d_M(\mu, g\mu)\big). \end{aligned}$$

□

Corollary 5.0.6 (L.B.C. property for reducible mapping classes)**.** *If $f, g \in \mathcal{MCG}(\mathcal{S})$ are conjugate reducible mapping classes of infinite order, then there is a conjugating element $\omega \in \mathcal{MCG}(\mathcal{S})$ with*

$$|\omega| \prec |f| + |g|.$$

Proof. Let $\mathcal{M}$ be the set of representative markings and let C be the constant bounding the diameter of $\mathcal{M}$. Let K be the constant depending only on $\mathcal{S}$ defined by

$$K = \max\{\ K_\mu \ \ : \ \ \mu \in \mathcal{M}\ \},$$

where K_μ is the constant associated to μ in Proposition 5.0.5. Suppose f_1 and f_2 are conjugate reducible mapping classes of infinite order. Let a_1 and a_2 be such that $a_i f_i a_i^{-1} = g_i$ have a good marking $\mu \in \mathcal{M}$, and satisfying $d_M(\mu_0, a_i\mu_0) \prec d_M(\mu_0, f_i\mu_0)$. Then each

$$d_M(\mu_0, g_i\mu_0) \leq 2d_M(\mu_0, a_i\mu_0) + d_M(\mu_0, f_i\mu_0) \prec d_M(\mu_0, f_i\mu_0).$$

By Proposition 5.0.5, there exists $\omega \in \mathcal{MCG}(\mathcal{S})$ such that $g_1\omega = \omega g_2$, and

$$\begin{aligned} d_M(\mu, \omega\mu) &\prec K_\mu\big(d_M(\mu, g_1\mu) + d_M(\mu, g_2\mu)\big) \\ &\leq K\big(d_M(\mu, g_1\mu) + d_M(\mu, g_2\mu)\big). \end{aligned}$$

Hence, by the triangle inequality,

$$\begin{aligned} d_M(\mu_0, \omega\mu_0) &\leq d_M(\mu_0, \mu) + d_M(\mu, \omega\mu) + d_M(\omega\mu, \omega\mu_0) \\ &\leq 2C + K d_M(\mu, g_1\mu) + d_M(\mu, g_2\mu)) \\ &\leq 2C + K\big(4C + d_M(\mu_0, g_1\mu_0) + d_M(\mu_0, g_2\mu_0)\big) \\ &\prec d_M(\mu_0, g_1\mu_0) + d_M(\mu_0, g_2\mu_0) \end{aligned}$$

Set $\omega' = a_1^{-1}\omega a_2$. Then ω' is a conjugator for f_1 and f_2, and

$$\begin{aligned} d_M(\mu_0, \omega'\mu_0) &\leq d_M(\mu_0, a_1\mu_0) + d_M(\mu_0, \omega\mu_0) + d_M(\mu_0, a_2\mu_0) \\ &\prec d_M(\mu_0, f_1\mu_0) + d_M(\mu_0, f_2\mu_0). \end{aligned}$$

□

References

[Ago10] Ian Agol, *Ideal triangulations of pseudo-Anosov mapping tori*, Preprint, 2010, `arXiv:1008.1606v2 [math.GT]`.

[BH95] M. Bestvina and M. Handel, *Train-tracks for surface homeomorphisms*, Topology **34** (1995), no. 1, 109–140.

[BH99] Martin R. Bridson and André Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 319, Springer-Verlag, Berlin, 1999.

[BKMM06] Jason Behrstock, Bruce Kleiner, Yair N. Minsky, and Lee Mosher, *Geometry and rigidity of mapping class groups*, Preprint, 2006, `arXiv:0801.2006 [math.GT]`.

[BM08] Jeffrey Brock and Howard Masur, *Coarse and synthetic Weil-Petersson geometry: quasi-flats, geodesics and relative hyperbolicity*, Geom. Topol. **12** (2008), no. 4, 2453–2495.

[Boo59] William W. Boone, *The word problem*, Ann. of Math. (2) **70** (1959), 207–265.

[Bow06] Brian H. Bowditch, *Intersection numbers and the hyperbolicity of the curve complex*, J. Reine Angew. Math. **598** (2006), 105–129.

[Deh11] M. Dehn, *Über unendliche diskontinuierliche gruppen*, Mathematische Annalen **71** (1911), no. 1, 116–144.

[ECH$^+$92] David B. A. Epstein, James W. Cannon, Derek F. Holt, Silvio V. F. Levy, Michael S. Paterson, and William P. Thurston, *Word processing in groups*, Jones and Bartlett Publishers, Boston, MA, 1992.

[FLP79] *Travaux de Thurston sur les surfaces*, Astérisque, vol. 66, Société Mathématique de France, Paris, 1979, Séminaire Orsay, With an English summary.

[FM08] Benson Farb and Dan Margalit, *A primer on mapping class groups*, Book in progres, 2008, `www.math.utah.edu/∼margalit/primer`.

[Gro87] M. Gromov, *Hyperbolic groups*, Essays in group theory, Math. Sci. Res. Inst. Publ., vol. 8, Springer, New York, 1987, pp. 75–263.

[Gro75] Edna K. Grossman, *On the residual finiteness of certain mapping class groups*, J. London Math. Soc. (2) **9** (1974/75), 160–164.

[Ham09] Ursula Hamenstädt, *Geometry of the mapping class group, II: A biautomatic structure*, Preprint, 2009, `arXiv:math/0912.0137v1 [math.GR]`.

[Har81] W. J. Harvey, *Boundary structure of the modular group*, Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978) (Princeton, N.J.), Ann. of Math. Stud., vol. 97, Princeton Univ. Press, 1981, pp. 245–251.

[Hem79] Geoffrey Hemion, *On the classification of homeomorphisms of* 2*-manifolds and the classification of* 3*-manifolds*, Acta Math. **142** (1979), no. 1-2, 123–155.

[HT80] A. Hatcher and W. Thurston, *A presentation for the mapping class group of a closed orientable surface*, Topology **19** (1980), no. 3, 221–237.

[Ker83] Steven P. Kerckhoff, *The Nielsen realization problem*, Ann. of Math. (2) **117** (1983), no. 2, 235–265.

[Lic64] W. B. R. Lickorish, *A finite set of generators for the homeotopy group of a 2-manifold*, Proc. Cambridge Philos. Soc. **60** (1964), 769–778. MR 0171269 (30 #1500)

[Lys89] Igor G. Lysënok, *Some algorithmic properties of hyperbolic groups*, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. 4, 814–832, 912.

[Man05] Jason Fox Manning, *Geometry of pseudocharacters*, Geom. Topol. **9** (2005), 1147–1185 (electronic).

[Min96] Yair N. Minsky, *A geometric approach to the complex of curves on a surface*, Topology and Teichmüller spaces (Katinkulta, 1995), World Sci. Publ., River Edge, NJ, 1996, pp. 149–158.

[Min02] ______, *The classification of Kleinian surface groups, I: Models and bounds*, Preprint, 2002, `arXiv:math/0302208v3 [math.GT]`.

[MM99] Howard A. Masur and Yair N. Minsky, *Geometry of the complex of curves. I. Hyperbolicity*, Invent. Math. **138** (1999), no. 1, 103–149.

[MM00] ______, *Geometry of the complex of curves. II. Hierarchical structure*, Geom. Funct. Anal. **10** (2000), no. 4, 902–974.

[Mos86] Lee Mosher, *The classification of pseudo-Anosovs*, Low-dimensional topology and Kleinian groups (Coventry/Durham, 1984), London Math. Soc. Lecture Note Ser., vol. 112, Cambridge Univ. Press, Cambridge, 1986, pp. 13–75.

[Mos95] ______, *Mapping class groups are automatic*, Ann. of Math. (2) **142** (1995), no. 2, 303–384.

[Mos03] ______, *Train track expansions of measured foliations*, Preprint, 2003, `andromeda.rutgers.edu/~mosher`.

[Mos07] ______, *MSRI Course on mapping class groups*, Lecture notes, 2007, `andromeda.rutger.edu/~mosher`.

[RS07] Kasra Rafi and Saul Schleimer, *Covers and the curve complex*, Preprint, 2007, `arXiv:math/0701719v2 [math.GT]`.

[Thu88] William P. Thurston, *On the geometry and dynamics of diffeomorphisms of surfaces*, Bull. Amer. Math. Soc. (N.S.) **19** (1988), no. 2, 417–431.